

# Recombinant Human CD3E&CD3D Protein

Catalog No.: RP00624 **Recombinant**

## Sequence Information

Species	Gene ID	Swiss Prot
Human	916&915	P07766(CD3E)&P04234(CD3D)

### Tags

C-hFc

### Synonyms

CD3; CD3e; CD3E; CD3d; T3D; CD3D; CD3E&CD3D; CD3 delta&CD3 epsilon

## Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE > 95% by HPLC

Calculated MW	Observed MW
37.8 kDa (CD3E) , 35.4 kDa (CD3D)	48-58 kDa (CD3E) , 42-48 kDa (CD3D)

### Endotoxin

< 1 EU/μg of the protein by LAL method

### Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

## Background

T-cell surface glycoprotein CD3 epsilon & CD3 delta chain, also known as CD3E & CD3D, are single-pass type I membrane proteins. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain.

## Basic Information

### Description

Recombinant Human CD3E&CD3D/CD3 epsilon&CD3 delta Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Asp23-Asp126(CD3E) & Phe22-Ala105(CD3D)) of Human CD3E&CD3D/CD3 epsilon&CD3 delta (Accession #P07766(CD3E)&P04234(CD3D)) fused with a C-hFc tag at the C-terminus.

### Bio-Activity

### Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.




| [www.abclonal.com.cn](http://www.abclonal.com.cn)

---


## Validation Data




Human CD3E&CD3D on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.




The purity of Human CD3E&CD3D was greater than 95% as determined by SEC-HPLC.



Immobilized Human CD3E&CD3D, hFc Tag at 0.5 µg/mL (100 µL/Well). Dose response curve for Biotinylated Anti-CD3E&CD3D Ab.1, hFc Tag with the EC<sub>50</sub> of 31.68ng/mL determined by ELISA.



Immobilized Human CD3E&CD3D, hFc Tag at 1 µg/mL (100ul/Well). Dose response curve for Anti-CD3E&CD3D Ab.2, mFc Tag with the EC<sub>50</sub> of 45.9ng/mL determined by ELISA (QC Test).



Immobilized Tebentafusp, His Tag at 1 µg/mL (100 µL/well) on the plate. Dose response curve for Human CD3E&CD3D, hFc Tag with the EC<sub>50</sub> of 8.7ng/mL determined by ELISA.