ABclonal

www.abclonal.com

Recombinant Human VCAM-1/CD106 Protein

Catalog No.: RP00972 Recombinant 1 Publications

Sequence Information

Species Gene ID Swiss Prot Human 7412 P19320

Tags C-His

Synonyms

CD106;INCAM-100;VCAM1

Product Information

SourcePurificationHEK293 cells≥ 95 % as

determined by SDS-PAGE.

PAGE

Calculated MW Observed MW

74.23 kDa 100-110 kDa

Endotoxin

Please contact us for more information.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

8	400-999-6126
<u> </u>	cn.market@abclonal.com.cn
<u>~</u>	www.abclonal.com.cn

Background

This protein also known as CD16, is a cell surface sialoglycoprotein belonging to the immunoglobulin superfamily. Two forms of VCAM-1 with either six or seven extracellular Ig-like domains are generated by alternative splicing, with the longer form predominant. VCAM-1 is an endothelial ligand for very late antigen-4 (VLA-4) and α4β7 integrin expressed on leukocytes, and thus mediates leukocyte-endothelial cell adhesion and signal transduction. VCAM-1 expression is induced on endothelial cells during inflammatory bowel disease, atherosclerosis, allograft rejection, infection, and asthmatic responses. During these responses, VCAM-1 forms a scaffold for leukocyte migration. VCAM-1 also activates signals within endothelial cells resulting in the opening of an "endothelial cell gate" through which leukocytes migrate. VCAM-1 has been identified as a potential anti-inflammatory therapeutic target, the hypothesis being that reduced expression of VCAM-1 will slow the development of atherosclerosis. In addition, VCAM-1-activated signals in endothelial cells are regulated by cytokines indicating that it is important to consider both endothelial cell adhesion molecule expression and function during inflammatory processes.

Basic Information

Description

Recombinant Human VCAM-1/CD106 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Phe25-Glu698) of human VCAM1 (Accession #NP 001069.1) fused with a 6×His tag at the C-terminus.

Bio-Activity

Measured by the ability of the immobilized protein to support the adhesion of U937 human histiocytic lymphoma cells. When 5 x 10E4 cells/well are added to human VCAM1 coated plates (10 μ g/mL with 100 μ L/well), approximately 80%-90% cells will adhere after 1 hour at 37°C.|2.Measured by its binding ability in a functional ELISA.Immobilized APC anti-human CD106 Antibody at 1 μ g/mL (25 μ L/well) can bind Human VCAM1 with a linear range of 0.46-16.9ng/mL.

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.

Validation Data



Recombinant Human VCAM-1/CD106 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized APC anti-human CD106 Antibody at 1 μ g/mL (25 μ L/well) can bind Human VCAM1 with a linear range of 0.46-16.9ng/mL.