

Recombinant Human IGF-I Protein

Catalog No.: RP00996 Recombinant 1 Publications

Sequence Information

Species **Gene ID** **Swiss Prot**
 Human 3479 P05019-1

Tags

C-hFc&His

Synonyms

IGF1;IGF-I;IGFI;MGF; IGFI; MGF

Background

Basic Information

Description

Recombinant Human IGF-I Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gly 49 - Ala 118) of human IGF1 (Accession #NP_001104755.1) fused with an Fc, 6×His tag at the C-terminus.

Bio-Activity

1. Measured by its ability to stimulate p70 S6 Kinase(Thr389) and p85 S6 Kinase(Thr412) autophosphorylation in 293T human embryonic kidney cells. 0.01-1 ng/mL of Recombinant Human IGF1 can effectively enhance p70 S6 Kinase(Thr389) and p85 S6 Kinase(Thr412) autophosphorylation. 2. Measured by its binding ability in a functional ELISA. Immobilized recombinant human IGFBP6 at 1 µg/mL (100 µL/well) can bind recombinant human IGF1 with a linear range of 30-250 ng/mL. 3. Measured in a cell proliferation assay using MCF-7 cells. The ED₅₀ for this effect is typically 7.5-30 ng/mL, corresponding to a specific activity of 3.33×10⁴-1.33×10⁵ units/mg.

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.

Product Information

Source

HEK293 cells

Purification

≥ 85 % as determined by SDS-PAGE.

Calculated MW

34.44 kDa

Observed MW

41 kDa

Endotoxin

< 0.1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.


Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.


Contact

	400-999-6126
	cn.market@abclonal.com.cn
	www.abclonal.com.cn


Validation Data




Recombinant Human IGF-I Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.




Recombinant Human IGF1 enhances p70S6 Kinase(Thr389) and p85 S6 Kinase(Thr412) autophosphorylation in 293T human embryonic kidney cells. 0.01-1ng of Recombinant Human IGF1 can effectively enhance p70S6 Kinase(Thr389) and p85 S6 Kinase(Thr412) autophosphorylation.



Measured by its binding ability in a functional ELISA. Immobilized recombinant human IGFBP6 at 1 μ g/mL (100 μ L/well) can bind recombinant human IGF1 with a linear range of 30-250ng/mL.



Recombinant Human IGF1 promotes the proliferation of MCF-7 cells. The ED₅₀ for this effect is typically 7.5-30 ng/mL.



Human kidney organoids were cultured with EGF(Cat. RP03287), FGF2(Cat. RP01042), FGF7(Cat. RP01717), FGF9(Cat. RP01710), FGF10(Cat. RP01140), IGF-(Cat. RP00996), NOG(Cat. RP01237), RSPO1(Cat. RP00071), WNT-3a(Cat. RP01618SLQ). And further, DKK-1(RP01343) was used to induce the establishment of cell polarity.