

Recombinant Human Transferrin/TF Protein

Catalog No.: RP00997 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	7018	P02787

Tags

C-His

Synonyms

HEL-S-71p;PRO1557;PRO2086;TFQTL1;TF;Transferrin

Product Information

Source	Purification
HEK293 cells	≥ 95 % as determined by SDS-PAGE; ≥ 95 % as determined by HPLC.

Calculated MW	Observed MW
76.02 kDa	76-90 kDa

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

Basic Information

Description

Recombinant Human Transferrin/TF Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Val20-Pro698) of human Transferrin (Accession #NP_001054.1) fused with a 6×His tag at the C-terminus.

Bio-Activity

1. Measured in a serum-free cell proliferation assay using MCF-7 human breast cancer cells. The ED₅₀ for this effect is 0.075-0.3 ng/mL. 2. Measured by its binding ability in a functional ELISA. Immobilized Recombinant human Transferrin at 2 μg/mL (100 μL/well) can bind Recombinant human TfR with a linear range of 4-17 ng/mL.

Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.

Contact


☎		400-999-6126
✉		cn.market@abclonal.com.cn




Validation Data




Recombinant Human Transferrin/TF Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



The purity of Human Transferrin Protein (Cat.RP00997) was greater than 95% as determined by SEC-HPLC.



Recombinant Human Transferrin promotes the proliferation of MCF-7 human breast cancer cells. The ED_{50} for this effect is 0.075-0.3 ng/mL.



Immobilized Recombinant human Transferrin at 2 μ g/mL (100 μ L/well) can bind Recombinant human Tfr with a linear range of 4-17 ng/mL.