

Recombinant Human Testican-1/SPOCK1 Protein

Catalog No.: RP01051 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	6695	Q08629

Tags

C-His

Synonyms

SPOCK;TIC1;TICN1;Testican-1;SPOCK1

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

Basic Information

Description

Recombinant Human Testican-1/SPOCK1 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Arg22-Trp439) of human Testican 1/SPOCK1 (Accession #NP_004589.1.) fused with a 6xHis tag at the C-terminus.

Bio-Activity

Measured by its ability to inhibit active Cathepsin L cleavage of a fluorogenic peptide substrate Z-LR-AMC. The IC50 value approximately is 1.7 nM.

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Contact

 | 400-999-6126


 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Validation Data



Recombinant Human Testican-1/SPOCK1 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 55-60kDa.



Recombinant human SPOCK1 inhibits active Cathepsin L cleavage of a fluorogenic peptide substrate Z-LR-AMC (Catalog#ES008). The IC50 value approximately is 1.7nM.