

Recombinant Mouse VEGF-A/VEGF164 Protein

Catalog No.: RP01060

Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
Mouse	22339	Q00731-2

Tags

N-His

Synonyms

MVCD1;VEGFA;VEGF;VPF;VEGFA \square 164 \square

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

Endotoxin

< 0.1 EU/ μ g of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μ m filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

Basic Information

Description

Recombinant Mouse VEGF-A/VEGF164 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Ala27-Arg190) of mouse VEGF 164 (Accession #NP_001273986.1.) fused with a 6 \times His tag at the N-terminus.

Bio-Activity

1. Measured by its binding ability in a functional ELISA. Immobilized Recombinant Mouse VEGF164 at 1 μ g/mL (100 μ L/well) can bind Recombinant Human VEGFR2 with a linear range of 8-30 ng/mL. 2. Measured in a cell proliferation assay using human umbilical vein endothelial cells (HUVEC). The ED₅₀ for this effect is typically 0.006-0.022 ng/mL, corresponding to a specific activity of 4.54 \times 10⁷-1.67 \times 10⁸ units/mg.

Storage

Store the lyophilized protein at -20 $^{\circ}$ C to -80 $^{\circ}$ C for long term.

After reconstitution, the protein solution is stable at -20 $^{\circ}$ C for 3 months, at 2-8 $^{\circ}$ C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data



Recombinant Mouse VEGF-A/VEGF164 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 25-30kDa.



Immobilized Recombinant Mouse VEGF164 at 1 µg/mL (100 µL/well) can bind Recombinant Human VEGFR2 with a linear range of 8-30 ng/mL.



Recombinant Mouse VEGF164 promotes the proliferation of human umbilical vein endothelial cells (HUVEC). The ED₅₀ for this effect is typically 0.006-0.022 ng/mL, corresponding to a specific activity of 4.54×10⁷-1.67×10⁸units/mg.