

Recombinant Mouse IFN-gamma Protein

Catalog No.: RP01070 Recombinant 6 Publications

Sequence Information

Species Gene ID Swiss ProtMouse 15978 P01580

Tags C-hFc&His

Synonyms

IFG;IFI;IFN gamma;Interferon Gamma;IFNG

Product Information

Source Purification HEK293 cells > 92% by SDS-PAGE.

Calculated MW Observed MW

44.70 kDa 55 kDa

Endotoxin

 $< 0.1 \; \text{EU/}\mu\text{g}$ of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

6	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Background

Interferon-gamma (IFN-gamma) is a secreted protein which belongs to the type II interferon family. IFN-gamma is produced by a variety of immune cells under inflammatory conditions, notably by T cells and NK cells. IFN-gamma, in addition to having antiviral activity, has important immunoregulatory functions. Interferon-gamma is a central regulator of the immune response and signals via the Janus Activated Kinase (JAK)-Signal Transducer and Activator of Transcription (STAT) pathway.

Basic Information

Description

Recombinant Mouse IFN-gamma Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Met1-Cys155) of mouse Interferon Gamma (Accession #NP_032363.1) fused with an Fc, 6×His tag at the Cterminus.

Bio-Activity

1.Measured by its binding ability in a functional ELISA. Immobilized recombinant Mouse IFNGR1 at 0.2 μ g/mL (100 μ L/well) can bind recombinant Mouse IFNG, the EC₅₀ of Mouse IFNG is 2.2 μ g/mL.|2.Recombinant Mouse IFN-gamma stimulates mouse bone marrow-derived macrophages and peritoneal macrophages differentiation into M1 Macrophages(classically activated macrophage). The expression of the related biological markers TNF-a, Nos2, IL-1 μ , and IL-6 was detected to be significantly upregulated.|


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20 $^{\circ}$ C for 3 months, at 2-8 $^{\circ}$ C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data




Recombinant Mouse IFN-gamma Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Recombinant Mouse IFN-gamma Protein is greater than 90% as determined by SEC-HPLC.



Immobilized recombinant Mouse IFNGR1 at $0.2\mu g/mL$ (100 $\mu L/well$) can bind recombinant Mouse IFNG,the EC $_{50}$ of Mouse IFNG is 2.2ng/mL.



M0 macrophages(Catalog # RP01216S) can be induced differentiation into M1 macrophages(CD86+CD206-) by Recombinant Mouse IFN-gamma Protein (50ng/mL) and LPS (100ng/mL).