

Recombinant Mouse Placenta growth factor/PIGF/PGF Protein

Catalog No.: RP01080 Recombinant

Sequence Information

Species Gene ID Swiss Prot Mouse 18654 P49764

Tags C-hFc&His

Synonyms

PGF;PLGF;PIGF2;PIGF;PGFL;SHGC-10760;

Product Information

Source Purification HEK293 cells ≥ 85 % as

determined by SDS-

PAGE.

Calculated MW Observed MW

44.66 kDa 60 kDa

Endotoxin

< 1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

a	400-999-6126
\bowtie	cn.market@abclonal.com.cn
$\overline{\mathfrak{S}}$	www.abclonal.com.cn

Background

Placenta growth factor (PIGF) is a member of the PDGF/VEGF family of growth factors that share a conserved pattern of eight cysteines. Alternative splicing results in at least three human mature PIGF forms containing 131 (PIGF-1), 152 (PIGF-2), and 203 (PIGF-3) amino acids (aa) respectively. Only PIGF-2 contains a highly basic heparinbinding 21 aa insert at the C-terminus. Human PIGF-1 shares 56%, 55%, 74% and 95% aa identity with the comparable isoform of mouse, rat, canine, and equine PIGF, respectively. PIGF is mainly found as variably glycosylated, secreted, 55-60 kDa disulfide linked homodimers. Mammalian cells expressing PIGF include villous trophoblasts, decidual cells, erythroblasts, keratinocytes, and some endothelial cells. Circulating PIGF increases during pregnancy, reaching a peak in mid-gestation; this increase is attenuated in preeclampsia. However, deletion of PIGF in the mouse does not affect development or reproduction. Postnatally, mice lacking PIGF show impaired angiogenesis in response to ischemia.

Basic Information

Description

Recombinant Mouse Placenta growth factor/PIGF/PGF Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Met1-Pro158) of mouse PLGF/PGF (Accession #NP_032853.1) fused with an Fc, 6×His tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Recombinant Human VEGFR1 at 500 ng/mL (100 μ L/well) can bind Recombinant Mouse PLGF with a linear range of 12-49 ng/mL.


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20 $^{\circ}$ C for 3 months, at 2-8 $^{\circ}$ C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Mouse Placenta growth factor/PIGF/PGF Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized Recombinant Human VEGFR1 at 500ng/mL (100 μ L/well) can bind Recombinant Mouse PLGF with a linear range of 12-49 ng/mL.