

Recombinant Human VEGF-A/VEGF165 Protein

Catalog No.: RP01150 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 7422 P15692-4

Tags

N-His

Synonyms

VEGFA; MVCD1; VEGF; VPF; vascular endothelial growth factor A;MVCD1;VEGF;VPF;L VEGFA;VEGF A

Product Information

Source

HEK293 cells

Purification ≥ 95 % as

determined by SDS-PAGE;≥ 95 % as determined by HPLC.

Calculated MW Observed MW

20.01 kDa 24-26 kDa

Endotoxin

< 0.1 EU/ μ g of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

2		400-999-6126
\bowtie	T	cn.market@abclonal.com.cn

Background

Basic Information

Description

Recombinant Human VEGF-A/VEGF165 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ala27-Arg191) of human VEGF165 (Accession $\#NP_001165097.1$) fused with a $6\times His$ tag at the N-terminus.

Bio-Activity

1.Measured by its binding ability in a functional ELISA. Immobilized Recombinant Human VEGF165 at 1 μ g/mL (100 μ L/well) can bind Recombinant Human VEGFR2 with a linear range of 8-20 ng/mL.|2.Measured by its binding ability in a functional ELISA. Immobilized Human VEGF165 at 2 μ g/mL (100 μ L/well) can bind Human KDR with a linear range of 0.2-11.6 ng/mL.|3.Recombinant Human VEGF165 stimulates cell proliferation of the human umbilical vein endothelial cells (HUVEC). The ED50 for this effect is typically 0.19-0.78 ng/mL, corresponding to a specific activity of 1.28 \times 106 \sim 5.26 \times 106 units/mg.

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Validation Data




Recombinant Human VEGF-A/VEGF165 Protein was determined by SDS-PAGE under reducing (R) and non-reducing (NR) conditions.




The purity of human VEGF165 Protein (Cat.RP01150) was greater than 95% as determined by SEC-HPLC.



Immobilized Recombinant Human VEGF165 at 1 μ g/mL (100 μ L/well) can bind Recombinant Human VEGFR2 with a linear range of 8-20 ng/mL.



Immobilized Recombinant Human VEGF165 at 2 μ g/mL (100 μ L/well) can bind Human KDR with a linear range of 0.2-11.6 ng/mL.



Recombinant Human VEGF165 stimulates cell proliferation of the human umbilical vein endothelial cells (HUVEC). The ED $_{50}$ for this effect is typically 0.19-0.78 ng/mL, corresponding to a specific activity of 1.28 × $10^6 \sim 5.26 \times 10^6$ units/mg.