

# Recombinant Human FLT-2/FGFR-1/CD331 Protein

Catalog No.: RP01151 **Recombinant**

## Sequence Information

Species	Gene ID	Swiss Prot
Human	2260	P11362-1

### Tags

C-hFc&His

### Synonyms

FGFR1;BFGFR;CD331;CEK;ECCL;FGFBR;FGFR-1;FLG;FLT-2;FLT2;HBGFR;HH2;HRTFDS;KAL2;N-SAM;OGD;bFGF-R-1

## Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

### Endotoxin

< 0.1 EU/μg of the protein by LAL method.

### Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Background

## Basic Information

### Description

Recombinant Human FLT-2/FGFR-1/CD331 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Arg22-Glu376) of human FGF R1/CD331 (Accession #NP\_075598.2) fused with an Fc, 6×His tag at the C-terminus.

### Bio-Activity

1. Measured by its binding ability in a functional ELISA. Immobilized Recombinant Human FGF1 at 10 μg/mL (100 μL/well) can bind Recombinant Human FGFR1 with a linear range of 1.1-3.4 μg/mL. 2. Measured by its ability to inhibit FGF-acidic dependent proliferation of Balb/c 3T3 mouse fibroblasts. The ED<sub>50</sub> for this effect is typically 0.605-2.42 ng/mL.

### Storage

Store the lyophilized protein at -20°C to -80 °C for long term.

After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.

Avoid repeated freeze/thaw cycles.


## Contact

☎ | 400-999-6126


✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)


Validation Data



Recombinant Human FLT-2/FGFR-1/CD331 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 100-110 kDa..



Immobilized Recombinant Human FGF1 at 10μg/mL (100 μL/well) can bind Recombinant Human FGFR1 with a linear range of 1.1-3.4 μg/mL.



Recombinant Human FGFR1 inhibits FGF-acidic dependent proliferation of Balb/c 3T3 mouse fibroblasts. The ED<sub>50</sub> for this effect is typically 0.605-2.42 ng/mL.