ABclonal www.abclonal.com

Recombinant Human TREM2 Protein

Catalog No.: RP01159 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 54209 Q9NZC2

Tags

C-His

Synonyms

PLOSL2; TREM-2; Trem2a; Trem2b; Trem2c;TREM2;TREM-2;Trem2a;Trem2b; Trem2c

Product Information

Source Purification
HEK293 cells > 95% by SDSPAGE.

Calculated MW Observed MW

18.27 kDa 25-39 kDa

Endotoxin

 $< 0.1 \; \text{EU/}\mu\text{g}$ of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

6	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Background

Basic Information

Description

Recombinant Human TREM2 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (His19-Ser174) of human TREM-2 (Accession $\#NP_061838$) fused with a $6\times His$ tag at the C-terminus.

Bio-Activity

1.Measured by its binding ability in a functional ELISA.Immobilized Human TREM2 at $1\mu g/mL$ (100 $\mu L/well$) can bind TREM2 Rabbit pAb with a linear range of 0.06-1.86 ng/mL.


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human TREM-2 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 25-39 kDa..



Immobilized Human TREM2 at $1\mu g/mL$ (100 $\mu L/well)$ can bind TREM2 Rabbit pAb with a linear range of 0.06-1.86 ng/mL.