

Recombinant Human/Mouse/Rat Irisin/FNDC5 Protein

Catalog No.: RP01164 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human/Mouse/Rat	252995	Q8NAU1

Tags
C-His

Synonyms
FRCP2;irisin;FNDC5

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

Calculated MW	Observed MW
13.43 kDa	20-28 kDa

Endotoxin
< 0.1 EU/μg of the protein by LAL method.

Formulation
Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution
Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Background

Basic Information

Description

Recombinant Human/Mouse/Rat Irisin/FNDC5 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Asp32-Glu143) of Human/Mouse/Rat Irisin (Accession #NP_715637.2) fused with a 6×His tag at the C-terminus.


Bio-Activity

Measured by its ability to induce p38 MAPK activation in 3T3 L1 mouse embryonic fibroblast adipose-like cells. 0.1 μg/mL of Recombinant Human Irisin can effectively induce p38 MAPK activation.


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.


Validation Data



Recombinant Human/Mouse/Rat Irisin/FNDC5 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



The purity of Human/Mouse/Rat Irisin Protein (Cat.RP01164) was greater than 90% as determined by SEC-HPLC.



3T3-L1 preadipocytes were incubated with 1 pg/mL-0.1μg/mL of Recombinant Human Irisin for 12 minutes. The cells were lysed and Western blots were performed with anti-phospho p38 MAPK and total p38 MAPK antibodies.