

# Recombinant Mouse Latent TGF-beta 1 Protein

Catalog No.: RP01167   **Recombinant**   **1 Publications**

## Sequence Information

Species	Gene ID	Swiss Prot
Mouse	21803	P04202

### Tags

N-His

### Synonyms

TGF-beta1;Tgfb;Tgfb-1;TGFbeta1;CED;DPD1;LAP;TGFB;TGFbeta;TGFB1

## Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

### Endotoxin

< 0.1 EU/μg of the protein by LAL method.

### Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile 4 mM HCl. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

## Background

### Basic Information

#### Description

Recombinant Mouse Latent TGF-beta 1 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Leu30-Ser390) of mouse TGF-beta 1 (Accession #NP\_035707.1) fused with a 8xHis tag at the N-terminus.

#### Bio-Activity

Measured by its ability to inhibit the IL-4-dependent proliferation of HT-2 mouse T cells. The ED<sub>50</sub> for this effect is 0.02-0.1 ng/mL, corresponding to a specific activity of 1.0×10<sup>7</sup>-5.0×10<sup>7</sup> units/mg.

#### Storage

Store the lyophilized protein at -20°C to -80 °C for long term.

After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.

Avoid repeated freeze/thaw cycles.


## Contact

 | 400-999-6126


 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

# Validation Data



Active Recombinant Mouse TGF-beta 1 Protein was determined by SDS-PAGE with Coomassie Blue, showing bands at 14 kDa, 40-50 kDa and 55 kDa..



Recombinant Mouse TGF-beta 1 inhibits the IL-4-dependent proliferation of HT-2 mouse T cells. The ED<sub>50</sub> for this effect is 0.02-0.1 ng/mL, corresponding to a specific activity of 1.0×10<sup>7</sup>-5.0×10<sup>7</sup>units/mg.