

Recombinant Human Endothelial protein C receptor/CD201 Protein

Catalog No.: RP01181 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 10544 Q9UNN8

Tags

C-His

Synonyms

PROCR;CCCA;CCD41;EPCR;h-PROCR

Product Information

Source

Purification

HEK293 cells

> 95% by SDS-PAGE.

Endotoxin

 $< 1.0 \; \text{EU/}\mu\text{g}$ of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

2	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Background

Basic Information

Description

Recombinant Human Endothelial protein C receptor/CD201 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ser18-Ser210) of human EPCR/CD201/PROCR (Accession #NP_006395.2) fused with a 6×His tag at the C-terminus.


Bio-Activity

Storage

Store the lyophilized protein at -20 $^{\circ}$ C to -80 $^{\circ}$ C for long term. After reconstitution, the protein solution is stable at -20 $^{\circ}$ C for 3 months, at 2-8 $^{\circ}$ C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human Endothelial protein C receptor/CD201 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 36-40 kDa.