

Recombinant Human Siglec-15/CD33L3 Protein

Catalog No.: RP01187 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	284266	Q6ZMC9

Tags

C-hFc&His

Synonyms

CD33L3; HsT1361;
SIGLEC-15;SIGLEC15;Siglec-15

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

Calculated MW	Observed MW
53.34 kDa	60-70 kDa

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS,300mM NaCl, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Background

Basic Information

Description

Recombinant Human Siglec-15/CD33L3 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Phe20-Thr263) of human Siglec-15/CD33L3 (Accession #NP_998767.1) fused with a Fc, 6×His tag at the C-terminus.

Bio-Activity

Measured by its ability to inhibit Anti-CD3-induced proliferation of jurkat cells. The ED₅₀ for this effect is 1.8-7.2 μg/mL.


Storage

Store at -20°C.Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data



Active Recombinant Human Siglec-15/CD33L3 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 60 kDa.



The purity of human Siglec-15/CD33L3 Protein (Cat.RP01187) was greater than 95% as determined by SEC-HPLC.



Recombinant Human Siglec-15 inhibits Anti-CD3-induced proliferation of jurkat cells. The ED_{50} for this effect is 1.8-7.2 $\mu\text{g/mL}$.