

# Recombinant Human IL-15RA/CD215 Protein

Catalog No.: RP01192 Recombinant

### **Sequence Information**

Species Gene ID Swiss Prot Human 3601 Q13261

Tags

C-hFc&His

**Synonyms** IL15RA;CD215

### **Product Information**

**Source**HEK293 cells
Purification
> 95% by SDSPAGE.

#### **Endotoxin**

< 0.1 EU/ $\mu$ g of the protein by LAL method.

#### **Formulation**

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

### **Contact**

<b>a</b>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

### **Background**

### **Basic Information**

#### Description

Recombinant Human IL-15RA/CD215 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Met1-Thr205) of human IL-15R alpha/CD215 (Accession #NP\_002180.1) fused with a Fc, 6×His tag at the C-terminus.

#### **Bio-Activity**


Measured by its binding ability in a functional ELISA. Immobilized recombinant Human IL15 at 2  $\mu$ g/mL (100  $\mu$ L/well) can bind recombinant Human IL-15R alpha, the EC<sub>50</sub> of Human IL-15R alpha is 178.9 ng/mL.

#### Storage


Store the lyophilized protein at -20  $^{\circ}$ C to -80  $^{\circ}$ C for long term. After reconstitution, the protein solution is stable at -20  $^{\circ}$ C for 3 months, at 2-8  $^{\circ}$ C for up to 1 week.

Avoid repeated freeze/thaw cycles.

## **Validation Data**



Recombinant Human IL-15RA/CD215 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 75-90 kDa.



Immobilized recombinant Human IL15 at  $2\mu g/mL$  ( $100\mu L/well$ ) can bind recombinant Human IL-15R alpha, the EC $_{50}$  of Human IL-15R alpha is 178.9ng/mL.