

Recombinant Human Serpin D1/Heparin cofactor 2 Protein

Catalog No.: RP01203 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 3053 P05546

Tags

C-His

Synonyms

SERPIND1;D22S673;HC2;HCF2;HCII;HLS2;LS2;THPH10

Product Information

Source Purification HEK293 cells > 90% by SDS-PAGE.

Calculated MW Observed MW

55.78 kDa 70-80 kDa

Endotoxin

< 0.1 EU/ μ g of the protein by LAL method

Formulation

Lyophilized from a 0.22 µm filtered solution of 50mM Tris,150mM NaCl, pH7.5.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

a	400-999-6126
×	cn.market@abclonal.com.cn
	www.abclonal.com.cn

Background

Basic Information

Description

Recombinant Human Serpin D1/Heparin cofactor 2 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gly20-Ser499) of human Serpin D1/Heparin Cofactor II (Accession $\#NP_000176.2$.) fused with a 6×His tag at the C-terminus.

Bio-Activity

Measured by its ability to inhibit Recombinant Human Coagulation Factor II/Thrombin cleavage of a fluorogenic peptide substrate Boc-VPR-AMC.The IC50 value is <0.19 nM.


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20 $^{\circ}$ C for 3 months, at 2-8 $^{\circ}$ C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human Serpin D1/Heparin cofactor 2 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 70-80 kDa.



Recombinant human SERPIND1 inhibits Recombinant Human Coagulation Factor II/Thrombin cleavage of a fluorogenic peptide substrate Boc-VPR-AMC.The IC50 value is <0.19 nM.