

Recombinant Mouse CSF-1/M-CSF Protein

Catalog No.: RP01216 **Recombinant** **6 Publications**

Sequence Information

Species	Gene ID	Swiss Prot
Mouse	12977	P07141-1

Tags

C-His

Synonyms

MCSF;M-CSF;CSF-1;Lanimostim;CSF1

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

Calculated MW	Observed MW
30.19 kDa	40-55 kDa

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

Macrophage colony-stimulating factor 1, also known as CSF-1, M-CSF, is a single-pass membrane protein which is disulfide-linked as a homodimer or heterodimer. Granulocyte / macrophage colony-stimulating factors are cytokines that act in hematopoiesis by controlling the production, differentiation, and function of 2 related white cell populations of the blood, the granulocytes and the monocytes-macrophages. M-CSF/CSF-1 is known to facilitate monocyte survival, monocyte-to-macrophage conversion, and macrophage proliferation. M-CSF/CSF-1 is a secreted cytokine which influences hemopoietic stem cells to differentiate into macrophages or other related cell types. It binds to the Colony stimulating factor 1 receptor. M-CSF/CSF-1 may also be involved in development of the placenta. The active form of M-CSF/CSF-1 is found extracellularly as a disulfide-linked homodimer, and is thought to be produced by proteolytic cleavage of membrane-bound precursors. M-CSF/CSF-1 induces cells of the monocyte/macrophage lineage. It also plays a role in immunological defenses, bone metabolism, lipoproteins clearance, fertility and pregnancy. Upregulation of M-CSF/CSF-1 in the infarcted myocardium may have an active role in healing not only through its effects on cells of monocyte/macrophage lineage, but also by regulating endothelial cell chemokine expression.

Basic Information

Description

Recombinant Mouse CSF-1/M-CSF Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Met1-Glu262) of mouse M-CSF/CSF-1 (Accession #NP_031804.3.) fused with a 6×His tag at the C-terminus.


Bio-Activity

1. Measured in a cell proliferation assay using M-NFS-60 mouse myelogenous leukemia lymphoblast cells. The ED₅₀ for this effect is typically 0.1-0.4 ng/mL, corresponding to a specific activity of 2.5×10⁵-1.0×10⁷ units/mg. | 2. Measured in a cell proliferation assay using mouse bone marrow cells. The ED₅₀ for this effect is 7.5-30.1 ng/mL, corresponding to a specific activity of 3.32×10⁴~1.33×10⁵ units/mg. | 3. Measured in a cell proliferation assay using M-NFS-60 mouse myelogenous leukemia lymphoblast cells. The ED₅₀ for this effect is 2-8 ng/mL, corresponding to a specific activity of 12.5×10⁴~5.0×10⁵ units/mg.


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.


Validation Data



Active Recombinant Mouse CSF-1/M-CSF Protein was determined by SDS-PAGE with Coomassie Blue



Recombinant Mouse CSF-1/M-CSF stimulates cell proliferation of the mouse bone marrow cells. The ED_{50} for this effect is 7.5-30.1 ng/mL, corresponding to a specific activity of $3.32 \times 10^4 \sim 1.33 \times 10^5$ units/mg.



Recombinant Mouse CSF-1/M-CSF stimulates cell proliferation of the M-NFS-60 mouse myelogenous leukemia lymphoblast cells. The ED_{50} for this effect is 8.31-33.24 ng/mL, corresponding to a specific activity of $3.01 \times 10^4 \sim 1.20 \times 10^5$ units/mg.