

Recombinant Human IL-15 Protein

Catalog No.: RP01236 Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
Human	3600	P40933-1

Tags

C-His

Synonyms

IL15;IL-15

Background

Basic Information

Description

Recombinant Human IL-15 Protein is produced by *E. coli* expression system. The target protein is expressed with sequence (Asn49-Ser162) of human IL-15 (Accession #NP_000576.1) fused with a 6×His tag at the C-terminus.

Bio-Activity

Measured in a cell proliferation assay using M07e human megakaryocytic leukemic cells. The ED₅₀ for this effect is 1.40-5.60 ng/mL, corresponding to a specific activity of $1.78 \times 10^5 \sim 7.14 \times 10^5$ units/mg.

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.

Product Information

Source

E. coli

Purification

≥ 95 % as determined by SDS-PAGE.

Calculated MW

13.61 kDa

Observed MW

15 kDa

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.


Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.


Contact

	400-999-6126
	cn.market@abclonal.com.cn
	www.abclonal.com.cn

Validation Data



Recombinant Human IL-15 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Measured in a cell proliferation assay using M07e human megakaryocytic leukemic cells. The ED₅₀ for this effect is 1.40-5.60 ng/mL, corresponding to a specific activity of $1.78 \times 10^5 \sim 7.14 \times 10^5$ units/mg.