Recombinant Human IL-3 Protein

www.abclonal.com

ABclonal

Catalog No.: RP01362 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 3562 P08700

Tags

C-His

Synonyms

IL3;IL-3;MCGF;MULTI-CSF

Product Information

Source Purification HEK293 cells ≥ 95 % as

determined by SDS-

PAGE.

Calculated MW Observed MW

15.93 kDa 20-35 kDa

Endotoxin

< 1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	Τ	www.abclonal.com.cn

Background

IL3 (interleukin 3), also known as IL-3, is a potent growth-promoting cytokine that belongs to the IL-3 family. IL3/IL-3 also belongs to the group of interleukins. Interleukins are produced by a wide variety of body cells. The function of the immune system depends in a large part on interleukins, and rare deficiencies of a number of them have been described, all featuring autoimmune diseases or immune deficiency. The majority of interleukins are synthesized by helper CD4+ T lymphocytes, as well as through monocytes, macrophages, and endothelial cells. They promote the development and differentiation of T, B, and hematopoietic cells. IL3/IL-3 is capable of supporting the proliferation of a broad range of hematopoietic cell types. It is involved in a variety of cell activities such as cell growth, differentiation, and apoptosis. IL3/IL-3 has been shown to also possess neurotrophic activity, and it may be associated with neurologic disorders.

Basic Information

Description

Recombinant Human IL-3 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ala20-Phe152) of human interleukin-3/IL-3 (Accession #NP 000579.2) fused with a 6×His tag at the C-terminus.

Bio-Activity

1.Measured in a cell proliferation assay using TF-1 human erythroleukemic cells. The ED $_{50}$ for this effect is 1.5-6 ng/mL.|2.Recombinant Human TPO(50 ng/mL, Cat. RP00174) , IL-3(15 ng/mL), IL-6(15 ng/mL, Cat. RP00201) and IL-11(15 ng/mL, Cat. RP00050) induce hematopoietic stem and progenitor cells to differentiate into megakaryocytes. After 6 days, the induction of CD41/42+ megakaryocytes was successful.

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20 $^{\circ}\text{C}$ for 3 months, at 2-8 $^{\circ}\text{C}$ for up to 1 week.

Avoid repeated freeze/thaw cycles.


Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.

Validation Data



Recombinant Human IL-3 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Recombinant Human IL3 promotes the proliferation of TF 1 human erythroleukemic cells. The ED $_{50}$ for this effect is 0.345-1.38 ng/mL, corresponding to a specific activity of $7.25 \times 10^5 \sim 2.90 \times 10^6$ units/mg.