

**Catalog No.: RP01365** **Recombinant**

Species	Gene ID	Swiss Prot
Human	131450	O8TD46

## C-hFc&amp;His

CD200R1:CD200R:HCRT2:MOX2R:OX2R

<b>Source</b>	<b>Purification</b>
HEK293 cells	≥ 95 % as determined by SDS-PAGE

53.59 kDa                      70-100 kDa

< 1 EU/μg of the protein by LAL method.

Lyophilized from a 0.22  $\mu\text{m}$  filtered solution of PBS, pH 7.4.

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

The cluster of differentiation (CD) system is commonly used as cell markers in Immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. Cell surface glycoprotein CD200 receptor 1 (CD200R1) is an isoform of CD200 receptors that is expressed on cells of the myeloid lineage. CD200R1 is a receptor for the OX-2 membrane glycoprotein. The receptor-substrate interaction may serve as a myeloid downregulatory signal.


Recombinant Human CD200R1 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ala27-Leu266) of human CD200R/MOX2R (Accession #NP\_620161.1) fused with a Fc, 6×His tag at the C-terminus.

Measured by its binding ability in a functional ELISA. Immobilized Human CD200 at 2 µg/mL (100 µL/well) can bind Human CD200 R with a linear range of 0.058-8.34 ng/mL.


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.  
After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.  
Avoid repeated freeze/thaw cycles.

 | 400-999-6126


 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

Validation Data



Recombinant Human CD200R1 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized Human CD200 at 2  $\mu$ g/mL (100  $\mu$ L/well) can bind Human CD200 R with a linear range of 0.058-8.34 ng/mL.