

Recombinant Human Fc-gamma RIII alpha/CD16a(F176V) Protein

Catalog No.: RP01366 **Recombinant**

Sequence Information

Species **Gene ID** **Swiss Prot**
Human 2214□F176V□P08637

Tags

C-His

Synonyms

CD16;CD16A;FCG3;FCGR3;FCGRIII;FCR-1
0;FCRIII;FCRIIIA;IGFR3;IMD20;FCGR3A

Product Information

Source **Purification**
HEK293 cells > 95% by SDS-
PAGE.

Endotoxin

<1EU/μg

Formulation

Lyophilized from a 0.22 μm filtered
solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening.
Reconstitute to a concentration of
0.1-0.5 mg/mL in sterile distilled water.
Avoid vortex or vigorously pipetting the
protein. For long term storage, it is
recommended to add a carrier protein or
stabilizer (e.g. 0.1% BSA, 5% HSA, 10%
FBS or 5% Trehalose), and aliquot the
reconstituted protein solution to
minimize free-thaw cycles.

Contact

☎ | 400-999-6126
✉ | cn.market@abclonal.com.cn
🌐 | www.abclonal.com.cn

Background

This protein is a receptor for the Fc portion of immunoglobulin G, and it is involved in the removal of antigen-antibody complexes from the circulation, as well as other other antibody-dependent responses. The protein is expressed on natural killer (NK) cells as an integral membrane glycoprotein anchored through a transmembrane peptide, whereas FCGR3B is expressed on polymorphonuclear neutrophils (PMN) where the receptor is anchored through a phosphatidylinositol (PI) linkage. Mutations in this gene have been linked to susceptibility to recurrent viral infections, susceptibility to systemic lupus erythematosus, and alloimmune neonatal neutropenia.

Basic Information

Description


Recombinant Human Fc-gamma RIII alpha/CD16a(F176V) Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gly17-Gln208(F176V)) of human FCGR3A/CD16A(F176V) (Accession #NP_001121065.1) fused with a 6×His tag at the C-terminus.

Bio-Activity

Storage

Store the lyophilized protein at -20°C to -80°C for long term.
After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.
Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human Fc-gamma RIII
alpha/CD16a(F176V) Protein was determined
by SDS-PAGE with Coomassie Blue, showing
a band at 38-52kDa.