# **Recombinant Human CCL21 Protein**



Catalog No.: RP01402 Recombinant

### **Sequence Information**

Species Gene ID Swiss Prot Human 6366 000585

Tags

No tag

**Synonyms** 

CCL21;6Ckine;CKb9;ECL;SCYA21;SLC;TC A4

### **Product Information**

Source Purification
E. coli > 85% by SDSPAGE.

Calculated MW Observed MW

12.25 kDa 16-17 kDa

### Endotoxin

Please contact us for more information.

#### **Formulation**

Lyophilized from a 0.22  $\mu$ m filtered solution of PBS, pH 7.4.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

#### **Contact**

<b>a</b>		400-999-6126
$\bowtie$	Τ	cn.market@abclonal.com.cn
<u>~</u>	Т	www.abclonal.com.cn

### **Background**

Chemokines are a family of small chemotactic cytokines, or proteins secreted by cells. Chemokines share the same structure similarities such as small size, and the presence of four cysteine residues in conserved locations in order to form their 3-dimensional shape. Some of the chemokines are considered pro-inflammatory which can be induced to recruit cells of the immune system to a site of infection during an immune response, while others are considered homeostatic and are implied in controlling the migration of cells during normal processes of tissue maintenance and development. There are four members of the chemokine family: C-C kemokines, C kemokines, CXC kemokines and CX3C kemokines. The C-C kemokines have two cysteines nearby the amino terminus. There have been at least 27 distinct members of this subgroup reported for mammals, called C-C chemokine ligands-1 to 28. Chemokine ligand 21(CCL21), also known as 6Ckine, exodus-2, and secondary lymphoid-tissue chemokine(SLC), is a small cytokine belonging to the C-C chemokine family. CCL21 takes its name 6Ckine for its consititutively six conserved cysteine residues but not four cysteines typical to chemokines. CCL21 has function in ininducing vigorous calcium migrations and chemotactic responses.

#### **Basic Information**

#### **Description**

Recombinant Human CCL21 Protein is produced by *E. coli* expression system. The target protein is expressed with sequence (Ser24-Pro134) of human CCL21/6Ckine (Accession #NP\_002980.1) fused with no additional amino acid.

#### **Bio-Activity**


#### **Storage**

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20  $^{\circ}$ C for 3 months, at 2-8  $^{\circ}$ C for up to 1 week.

Avoid repeated freeze/thaw cycles.

## **Validation Data**



Recombinant Human CCL21 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 16-17kDa.