

# Recombinant Human Endoglin/ENG/CD105 Protein

Catalog No.: RP01413 Recombinant

# **Sequence Information**

Species Gene ID Swiss Prot Human 2022 P17813

**Tags** C-His

Synonyms

END;HHT1;ORW1;CD105;ENG;endoglin

# **Product Information**

**Source** Purification HEK293 cells ≥ 95 % as

≥ 95 % as determined by SDS-

PAGE.

Calculated MW Observed MW

61.59 kDa 75-80 kDa

#### **Endotoxin**

< 0.1 EU/ $\mu g$  of the protein by LAL method.

## **Formulation**

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

#### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact

<b>a</b>	400-999-6126
×	cn.market@abclonal.com.cn
<b>⊙</b>	www.abclonal.com.cn

# **Background**

Endoglin, also known as CD105, is a type I homodimeric transmembrane glycoprotein with a large, disulfide-linked, extracellular region and a short, constitutively phosphorylated cytoplasmic tail. Endoglin contains an RGD tripeptide which is a key recognition structure in cellular adhesion, suggesting a critical role for endoglin in the binding of endothelial cells to integrins and/or other RGD receptors. Endoglin is highly expressed on vascular endothelial cells, chondrocytes, and syncytiotrophoblasts of term placenta. It is also found on activated monocytes, mesenchymal stem cells and leukemic cells of lymphoid and myeloid lineages. As an accessory receptor for the TGF- $\beta$  superfamily ligands, endoglin binds TGF- $\beta$ 1 and TGF- $\beta$ 3 with high affinity not by itself but by associating with TGF- $\beta$  type II receptor (T $\beta$ RII) and activates the downstream signal pathways. In addition, in human umbilical vein endothelial cells, ALK-1 is also a receptor kinase for endoglin threonine phosphorylation, and mutations in either of the two genes result in the autosomal-dominant vascular dysplasia, hereditary hemorrhagic telangiectasia (HHT). Endoglin has been regarded as a powerful biomarker of neovascularization, and is associated with several solid tumor types.

## **Basic Information**

### Description

Recombinant Human Endoglin/ENG/CD105 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Glu26-Gly586) of human Endoglin/CD105 (Accession #NP\_001108225.1) fused with a 6×His tag at the C-terminus.

## **Bio-Activity**

1.Measured by its binding ability in a functional ELISA. Immobilized Human CD105 at 2  $\mu g/mL$  (100  $\mu L/well)$  can bind Human ACVR2B with a linear range of 0.01-1.35  $\mu g/mL.|2.$ Measured by its binding ability in a functional ELISA. Immobilized Human CD105 at 2  $\mu g/mL$  (100  $\mu L/well$ ) can bind Human TGFBR2 with a linear range of 0.01-1.72  $\mu g/mL$ .

## **Shipping**

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

# Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


### **Operational Notes**

For your safety and health, please wear a lab coat and disposable gloves for handling.


# **Validation Data**



Recombinant Human Endoglin/ENG/CD105 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized Human CD105 at 2 $\mu$ g/mL (100  $\mu$ L/well) can bind Human ACVR2B with a linear range of 0.01-1.35 $\mu$ g/mL.



Immobilized Human CD105 at 2 $\mu$ g/mL (100  $\mu$ L/well) can bind Human TGFBR2 with a linear range of 0.01-1.72 $\mu$ g/mL.