

Biotinylated Recombinant Human MUC-16/CA125 Protein

Catalog No.: RP01417B Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 94025 Q8WXI7

Tags C-His&Avi

Synonyms

CA125; CA-125; CA125MUC-16; FLJ14303; MUC16

Product Information

Source Purification
HEK293 cells > 95% by Tris-Bis
PAGE

Calculated MW Observed MW

Endotoxin

< 1 EU/ μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	Τ	www.abclonal.com.cn

Background

MUC16, also known as the CA125 antigen, is a mucin protein that may be found in type I transmembrane or secreted forms that are used monitor the progress of epithelial ovarian cancer therapy. Thought to provide a protective, lubricating barrier against particles and infectious agents at mucosal surfaces. Binding to MSLN mediates heterotypic cell adhesion. This may contribute to the metastasis of ovarian cancer to the peritoneum by initiating cell attachment to the mesothelial epithelium via binding to MSLN.

Basic Information

Description

Biotinylated Recombinant Human MUC-16/CA125 Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Gly12660-Met12923) of Human MUC-16/CA125 fused with a His and Avi tag at the C-terminal.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80°C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Biotinylated Human CA125 on Tris-Bis PAGE under reduced conditions. The purity is greater than 95%.



Immobilized Human MSLN, hFc Tag at $0.5\mu g/ml$ (100 μ l/well) on the plate. Dose response curve for Biotinylated Human CA125, His Tag with the EC $_{50}$ of $0.35\mu g/ml$ determined by ELISA.