

Active Recombinant Human Mature TGF-beta 1 Protein

Catalog No.: RP01458

Recombinant

2 Publications

Sequence Information

Species	Gene ID	Swiss Prot
Human	7040	P01137

Tags

No tag

Synonyms

TGFB1; CED; DPD1; LAP; TGFB; TGFBeta; transforming growth factor beta-1; TGF-beta 1; CED; DPD1; LAP; TGFB; TGFBeta; TGF-β

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

Endotoxin

<0.1EU/μg

Formulation

Lyophilized from a 0.22 μm filtered solution of 50 mM Glycine, 150 mM NaCl, pH 3.5.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

TGF-beta 1 is a member of the transforming growth factor beta (TGF-beta) family. The transforming growth factor-beta family of polypeptides are involved in the regulation of cellular processes, including cell division, differentiation, motility, adhesion and death. TGF-beta 1 positively and negatively regulates many other growth factors. It inhibits the secretion and activity of many other cytokines including interferon-γ, tumor necrosis factor-alpha and various interleukins. It can also decrease the expression levels of cytokine receptors. Meanwhile, TGF-beta 1 also increases the expression of certain cytokines in T cells and promotes their proliferation, particularly if the cells are immature. TGF-beta 1 also inhibits proliferation and stimulates apoptosis of B cells, and plays a role in controlling the expression of antibody, transferrin and MHC class II proteins on immature and mature B cells. As for myeloid cells, TGF-beta 1 can inhibit their proliferation and prevent their production of reactive oxygen and nitrogen intermediates. However, as with other cell types, TGF-beta 1 also has the opposite effect on cells of myeloid origin. TGF-beta 1 is a multifunctional protein that controls proliferation, differentiation and other functions in many cell types. It plays an important role in bone remodeling as it is a potent stimulator of osteoblastic bone formation, causing chemotaxis, proliferation and differentiation in committed osteoblasts. Once cells lose their sensitivity to TGF-beta1-mediated growth inhibition, autocrine TGF-beta signaling can promote tumorigenesis. Elevated levels of TGF-beta1 are often observed in advanced carcinomas, and have been correlated with increased tumor invasiveness and disease progression.

Basic Information

Description

Active Recombinant Human Mature TGF-beta 1 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ala279-Ser390) of human Mature TGF-beta 1 (Accession #NP_000651.3) fused with no tag.

Bio-Activity

1. Measured by its binding ability in a functional ELISA. Immobilized Human TGF-beta 1 at 2 μg/mL (100 μL/well) can bind Human TGFB2 with a linear range of 0.78-11 ng/mL. 2. Measured by its ability to inhibit the IL-4 (Catalog: RP01161) dependent proliferation of HT-2 mouse T cells. The ED₅₀ for this effect is 42-168 pg/mL, corresponding to a specific activity of 6×10⁶~2.4×10⁷ units/mg.


Storage

Store the lyophilized protein at -20°C to -80°C for long term.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data




Active Recombinant Human Mature TGF-beta 1 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 14kDa.




Immobilized Human TGF-beta 1 at 2 µg/mL (100 µL/well) can bind Human TGFBR2 with a linear range of 0.78-11 ng/mL.




Recombinant Human Mature TGFβ1 inhibit the IL-4(Catalog: RP01161)-dependent proliferation of HT-2 mouse T cells. The ED₅₀ for this effect is 42-168 pg/mL,corresponding to a specific activity of 6×10⁶~2.4×10⁷ units/mg.




Active Recombinant Human mature TGF-beta1 induces EMT in A549 cells. As the concentration of TGF-beta1 increased, the expression of epithelial cell marker E-cad significantly downregulated.(Customer feedback data)



Active Recombinant Human mature TGF-beta1 stimulates smad2/3 autophosphorylation in Human Umbilical Vein Endothelial Cells (HUVEC). 10ng/mL of Recombinant Human TGF-beta 1 can effectively enhance smad2/3 autophosphorylation.(Customer feedback data)



Active Recombinant Human Mature TGF-beta 1 induces MCF7 cells EMT with a concentration of 12.5 ng/mL. After 10 h, the cell morphology significantly altered, appearing a fusiform shape (figure 1). After 48 h, the change of EMT associated genes are as figure 2. Results showed that the induction was successful.(Customer feedback data)



Recombinant mouse IL-23(Cat. RP02928LQ) and human TGF-β1 were used to induce the mouse T cells differentiate into Th17 cells. Flow cytometry was used to detect the expression of IL-17A. Results showed that Th17 cells were induced successfully.(Customer Feedback Data)