

Recombinant Human Glutathione S-transferase omega-1/GSTO-1/SPG-R Protein

Catalog No.: RP01494LQ Recombinant

Sequence Information

Species Gene ID **Swiss Prot** Human 9446 P78417

Tags

C-His

Synonyms

GSTO 1-1; GSTTLp28; HEL-S-21; P28; SPG-R;GSTO1; GSTO 1-1; GSTO1-1; glutathione S-transferase omega-1;GSTTLp28;HEL-S-21;P28;SPG-R;GSTO1-1

Product Information

Source

Purification E. coli

≥ 95 % as determined by SDS-PAGE.

Calculated MW Observed MW

28.27 kDa 30-35 kDa

Endotoxin

< 0.1 EU/ μg of the protein by LAL method.

Formulation

Supplied as a 0.22 µm filtered solution in 20 mM Tris,150 mM NaCl,10% glycerol,pH8.0.

Reconstitution

Background

Exhibits glutathione-dependent thiol transferase and dehydroascorbate reductase activities. Has S-(phenacyl)glutathione reductase activity. Has also glutathione Stransferase activity. Participates in the biotransformation of inorganic arsenic and reduces monomethylarsonic acid (MMA) and dimethylarsonic acid.

Basic Information

Description

Recombinant Human Glutathione S-transferase omega-1/GSTO-1/SPG-R Protein is produced by *E. coli* expression system. The target protein is expressed with sequence (Ser2-Leu241) of human GSTO1 (Accession #NP_004823.1) fused with a 6×His tag at the C-terminus.

Bio-Activity

1.Measured by its binding ability in a functional ELISA. Immobilized Human GSTO1 at 1 μg/mL (100 μL/well) can bind GSTO1 Rabbit mAb with a linear range of 0.5-4.33 ng/mL.|2.Measured by its binding ability in a functional ELISA. Immobilized Human Annexin A5 at 10 µg/mL (100 µL/well) can bind Human GSTO1 with a linear range of 0.15-3.16 µg/mL.


Storage

Store at -70°C. This product is stable at \leq -70°C for up to 1 year from the date of receipt. For optimal storage, aliquot into smaller quantities after centrifugation and store at recommended temperature. Avoid repeated freeze-thaw cycles. Avoid repeated freeze/thaw cycles.


Contact

6	400-999-6126
×<	cn.market@abclonal.com.cn
©	www.abclonal.com.cn


Validation Data



Recombinant Human Glutathione Stransferase omega-1/GSTO-1/SPG-R Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized recombinant Human GST01 at 1 μ g/mL (100 μ L/well) can bind GST01 Rabbit mAb with a linear range of 0.5-4.33ng/mL.



Immobilized recombinant Human Annexin A5 at $10\mu g/mL$ ($100~\mu L/well$) can bind Human GSTO1 with a linear range of 0.15-3.16 $\mu g/mL$.