

Catalog No.: RP01516LQ **Recombinant**

Species	Gene ID	Swiss Prot
Mouse	100043314	P86176

C-mFc

Synonyms
TIGIT;VSIG9;VSTM3;TIGIT;TIGIT;VSIG9;VS
TM3;TIGIT

Source	Purification
HEK293 cells	≥ 90% as determined by SDS-PAGE.

38.32 kDa 45-60 kDa

< 0.1 EU/μg of the protein by LAL method.

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water.

TIGIT, also known as V-set and transmembrane domain-containing protein 3 (VSTM3) or V-set and immunoglobulin domain-containing protein 9 (VSIG9) is a new surface protein containing an immunoglobulin variable domain, a transmembrane domain and an immunoreceptor tyrosine-based inhibitory motif (ITIM). TIGIT is expressed on regulatory, memory, activated T cells and NK cells. It binds PVR with high affinity, and PVRL2 with lower affinity, but not PVRL3. Knockdown of TIGIT with siRNA in human memory T cells did not affect T cell responses, however, TIGIT inhibits NK cytotoxicity directly through its ITIM. TIGIT suppresses T cell activation by promoting the generation of mature immunoregulatory dendritic cells. The binding of PVR to TIGIT on human dendritic cells enhanced the production of IL-1 and diminished the production of IL-12p4. Also, TIGIT counter inhibits the NK-mediated killing of tumor cells and protects normal cells from NK-mediated cytotoxicity thus providing an "alternative self" mechanism for MHC class I inhibition.

Recombinant Human TIGIT Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gly26-Gly141) of human TIGIT (Accession #NP_001139797.1) fused with a mFc tag at the C-terminus.

Measured by its binding ability in a functional ELISA. Immobilized Mouse CD155 at 2 $\mu\text{g/mL}$ (100 $\mu\text{L/well}$) can bind Mouse TIGIT with a linear range of 2-9 ng/mL.


Store at -70°C. This product is stable at $\leq -70^{\circ}\text{C}$ for up to 1 year from the date of receipt. For optimal storage, aliquot into smaller quantities after centrifugation and store at recommended temperature. Avoid repeated freeze-thaw cycles.

 | 400-999-6126


 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Validation Data



Recombinant Mouse TIGIT Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized Recombinant Mouse CD155 at 2 μ g/mL (100 μ L/well) can bind Mouse TIGIT with a linear range of 2-9 ng/mL.