Leader in Biomolecular Solutions for Life Science

www.abclonal.com

Recombinant Mouse c-kit/CD117 Protein

Catalog No.: RP01519 Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
Mouse	16590	P05532

Tags
C-Rabbit Fc

Synonyms
W; Bs; Fdc; Ssm; SCO1; SCO5; SOW3; CD117; c-KIT; Tr-kit; Gsfsco1; Gsfsco5; Gsfsow3;CD117

Product Information

Source	Purification
HEK293 cells	≥ 95 % as determined by SDS-PAGE.

Calculated MW	Observed MW
81.13 kDa	90-110 kDa

Endotoxin
< 0.1 EU/μg of the protein by LAL method.

Formulation
Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution
Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

* □□□□□□□□□□□□□□□□□□□□□□□□□□□□□

Background

C-Kit is a type 3 transmembrane receptor for MGF (mast cell growth factor, also known as stem cell factor). c-Kit contains 5 Ig-like C2-type (immunoglobulin-like) domains and 1 protein kinase domain. It belongs to the protein kinase superfamily, tyr protein kinase family, and CSF-1/PDGF receptor subfamily. C-Kit contains 5 Ig-like C2-type (immunoglobulin-like) domains and 1 protein kinase domain. C-Kit has tyrosine-protein kinase activity. Binding of the ligands leads to the autophosphorylation of KIT and its association with substrates such as phosphatidylinositol 3-kinase. Antibodies to c-Kit are widely used in immunohistochemistry to help distinguish particular types of tumor in histological tissue sections. It is used primarily in the diagnosis of GISTs. In GISTs, c-Kit staining is typically cytoplasmic, with stronger accentuation along the cell membranes. C-Kit antibodies can also be used in the diagnosis of mast cell tumors and in distinguishing seminomas from embryonal carcinomas. Mutations in the c-Kit gene are associated with gastrointestinal stromal tumors, mast cell disease, acute myelogenous leukemia, and piebaldism. Defects in KIT are a cause of acute myelogenous leukemia (AML). AML is a malignant disease in which hematopoietic precursors are arrested in an early stage of development. Note=Somatic mutations that lead to constitutive activation of KIT are detected in AML patients.

Basic Information

Description
Recombinant Mouse CD117/c-kit Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ser25-Thr523) of mouse CD117/c-kit (Accession #NP_001116205.1) fused with a raFc tag at the C-terminus.

Bio-Activity
Measured by its binding ability in a functional ELISA.Immobilized Mouse CD117 (Catalog: RP01519) at 5 μg/mL (100 μL/well) can bind Mouse SCF (Catalog: RP01055) with a linear range of 0.01-2 ng/mL.

Shipping
Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage
Store at -20℃.Store the lyophilized protein at -20℃ to -80 ℃ up to 1 year from the date of receipt.
After reconstitution, the protein solution is stable at -20℃ for 3 months, at 2-8℃ for up to 1 week.
Avoid repeated freeze/thaw cycles.

Operational Notes
For your safety and health, please wear a lab coat and disposable gloves for handling.

Validation Data



Recombinant Mouse c-kit/CD117 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized Mouse CD117 (Catalog: RP01519) at 5 μg/mL (100 μL/well) can bind Mouse SCF (Catalog: RP01055) with a linear range of 0.01-2 ng/mL.