

# Recombinant Mouse c-kit/CD117 Protein

Catalog No.: RP01519 **Recombinant**

## Sequence Information

Species	Gene ID	Swiss Prot
Mouse	16590	P05532

### Tags

C-Rabbit Fc

### Synonyms

W; Bs; Fdc; Ssm; SCO1; SCO5; SOW3; CD117; c-KIT; Tr-kit; Gsfsc01; Gsfsc05; Gsfscow3; CD117

## Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

### Endotoxin

<0.1EU/ $\mu$ g

### Formulation

Lyophilized from a 0.22  $\mu$ m filtered solution of PBS, pH 7.4.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

C-Kit is a type 3 transmembrane receptor for MGF (mast cell growth factor, also known as stem cell factor). c-Kit contains 5 Ig-like C2-type (immunoglobulin-like) domains and 1 protein kinase domain. It belongs to the protein kinase superfamily, tyr protein kinase family, and CSF-1/PDGF receptor subfamily. C-Kit contains 5 Ig-like C2-type (immunoglobulin-like) domains and 1 protein kinase domain. C-Kit has tyrosine-protein kinase activity. Binding of the ligands leads to the autophosphorylation of KIT and its association with substrates such as phosphatidylinositol 3-kinase. Antibodies to c-Kit are widely used in immunohistochemistry to help distinguish particular types of tumor in histological tissue sections. It is used primarily in the diagnosis of GISTs. In GISTs, c-Kit staining is typically cytoplasmic, with stronger accentuation along the cell membranes. C-Kit antibodies can also be used in the diagnosis of mast cell tumors and in distinguishing seminomas from embryonal carcinomas. Mutations in the c-Kit gene are associated with gastrointestinal stromal tumors, mast cell disease, acute myelogenous leukemia, and piebaldism. Defects in KIT are a cause of acute myelogenous leukemia (AML). AML is a malignant disease in which hematopoietic precursors are arrested in an early stage of development. Note=Somatic mutations that lead to constitutive activation of KIT are detected in AML patients.

## Basic Information

### Description

Recombinant Mouse CD117/c-kit Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ser25-Thr523) of mouse CD117/c-kit (Accession #NP\_001116205.1) fused with a raFc tag at the C-terminus.


### Bio-Activity

### Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.

## Validation Data

---



Recombinant Mouse CD117/c-kit Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 100-110kDa.