

Recombinant Human CLEC7A/Dectin-1/CD369 Protein

Catalog No.: RP01523 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 64581 Q9BXN2

Tags

N-Rabbit Fc

Synonyms

CLEC7A;BGR;CANDF4;CLECSF12;DECTIN 1:SCARE2:CD369

Product Information

Source Purification HEK293 cells ≥ 95 % as

determined by SDS-PAGE.

Calculated MW Observed MW

46.19 kDa 70-80 kDa

Endotoxin

< 0.1 EU/ μ g of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

6	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Background

Dectin-1 was recently identified as the most important receptor for beta-glucan. It is a type II transmembrane protein which binds beta-1,3 and beta-1,6 glucans, and is expressed on most cells of the innate immune system and has been implicated in phagocytosis as well as killing of fungi by macrophages, neutrophils and dendritic cells. Recognition of beta-glucan by dectin-1 triggers effective immune response, including phagocytosis and proinflammatory factor production, to eliminate infecting fungi, which especially benefits immunocompromised patients against opportunistic fungal infection. In addition, dectin-1 is involved in the adaptive immune response as well as autoimmune diseases and immune tolerance. Dectin-1 can recognize and respond to live fungal pathogens and is being increasingly appreciated as having a key role in the innate responses to these pathogens. In addition to its exogenous ligands, Dectin-1 can recognize an unidentified endogenous ligand on T cells and may act as a co-stimulatory molecule. Recent studies have highlighted the importance of Dectin-1 in anti-fungal immunity, in both mice and humans, and have suggested a possible involvement of this receptor in the control of mycobacterial infections.

Basic Information

Description

Recombinant Human CLEC7A/Dectin-1/CD369 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Thr66-Met247) of human CLEC7A/Dectin-1/CD369 (Accession #NP_922938.1) fused with a raFc tag at the N-terminus.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human CLEC7A/Dectin-1/CD369 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.