

Recombinant Mouse B7-DC/PD-L2/CD273 Protein

Catalog No.: RP01585 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Mouse	58205	Q9WUL5

Tags

C-hFc

Synonyms

B7-DC;B7DC;bA574F11.2;Btdc;CD273;PD-L2;PDCD1L2;PDL2;PDCD1LG2

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

Endotoxin

<0.1EU/μg

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Background

Programmed Death Ligand 2 (PD-L2), also known as B7-DC and butyrophilin-like protein, is a member of the B7 family of proteins that provide signals for regulating T-cell activation and tolerance. PD-L2 is expressed on dendritic cells, subsets of activated CD4+ and CD8+ T cells, and memory B cells that differentiate into plasma cells. At inflammatory sites such as rheumatoid arthritis, allergen exposure, and virus infection, PD-L2 is up-regulated on synoviocytes, infiltrating macrophages, dendritic cells, and airway epithelial cells. PD-L2, along with B7-H1/PD-L1, binds to T cell PD-1 where it promotes IFN-gamma production and CD40 Ligand up-regulation while inhibiting IL-4 production. In addition, PD-L2 binds to RGM-B on macrophages and alveolar epithelial cells, supporting respiratory immune tolerance. In asthma, PD-L2 suppresses IL-5 and IL-13 production, promotes IL-12 production by dendritic cells, and supports allergen-induced airway hyper-responsiveness and mucus production.

Basic Information

Description

Recombinant Mouse B7-DC/PD-L2/CD273 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Leu20-Arg219) of mouse B7-DC/PD-L2/CD273 (Accession #NP_067371.1) fused with a hFc tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Mouse PD-L2 (Catalog: RP01585) at 10 μg/mL (100 μL/well) can bind Biotinylated Mouse PD-1 with a linear range of 19.5-622 ng/mL.


Storage

Store the lyophilized protein at -20°C to -80°C for 12 months.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Mouse B7-DC/PD-L2/CD273
Protein was determined by SDS-PAGE with
Coomassie Blue, showing a band at 64 kDa.



Immobilized Mouse PD-L2 (Catalog:
RP01585) at 10 µg/mL (100 µL/well) can bind
Biotinylated Mouse PD-1 with a linear range
of 19.5-622 ng/mL.