

Recombinant Mouse B7-DC/PD-L2/CD273 Protein

Catalog No.: RP01585 Recombinant

Sequence Information

Species Gene ID Swiss ProtMouse 58205 Q9WUL5

Tags

C-hFc

Synonyms

B7-

DC;B7DC;bA574F11.2;Btdc;CD273;PD-L2;PDCD1L2;PDL2;PDCD1LG2

Product Information

Source Purification HEK293 cells > 95% by SDS-PAGE.

Calculated MW Observed MW

48.5 kDa 60-75 kDa

Endotoxin

<0.1EU/µg

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

2		400-999-6126
\bowtie	<u>cn.marke</u>	t@abclonal.com.cn
•	ww	w.abclonal.com.cn

Background

Programmed Death Ligand 2 (PD-L2), also known as B7-DC and butyrophilin-like protein, is a member of the B7 family of proteins that provide signals for regulating T-cell activation and tolerance .PD-L2 is expressed on dendritic cells, subsets of activated CD4+ and CD8+ T cells, and memory B cells that differentiate into plasma cells. At inflammatory sites such as rheumatoid arthritis, allergen exposure, and virus infection, PD-L2 is up-regulated on synoviocytes, infiltrating macrophages, dendritic cells, and airway epithelial cells. PD-L2, along with B7-H1/PD-L1, binds to T cell PD-1 where it promotes IFN-gamma production and CD40 Ligand up-regulation while inhibiting IL-4 production. In addition, PD-L2 binds to RGM-B on macrophages and alveolar epithelial cells, supporting respiratory immune tolerance. In asthma, PD-L2 suppresses IL-5 and IL-13 production, promotes IL-12 production by dendritic cells, and supports allergen-induced airway hyper-responsiveness and mucus production.

Basic Information

Description

Recombinant Mouse B7-DC/PD-L2/CD273 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Leu20-Arg219) of mouse B7-DC/PD-L2/CD273 (Accession #NP_067371.1) fused with a hFc tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Mouse PD-L2 (Catalog: RP01585) at 10 μ g/mL (100 μ L/well) can bind Biotinylated Mouse PD-1 with a linear range of 19.5-622 ng/mL.


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Mouse B7-DC/PD-L2/CD273 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 64 kDa.



Immobilized Mouse PD-L2 (Catalog: RP01585) at 10 μ g/mL (100 μ L/well) can bind Biotinylated Mouse PD-1 with a linear range of 19.5-622 μ m/mL.