

Recombinant Human CCL16/HCC-4 Protein

Catalog No.: RP01595 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	6360	O15467

Tags
C-6*His

Synonyms

CCL16; IL16; NCC4; SCYA16; C-C motif chemokine 16; Chemokine CC-4; HCC-4; Chemokine LEC; IL-10-inducible chemokine; LCC-1; Liver-expressed chemokine; Lymphocyte and monocyte chemoattractant; LMC; Monotactin-1; MTN-1; NCC-4; Small-inducible cytokine A16

Product Information

Source	Purification
HEK293 cells	PBS

Endotoxin
<0.01EU/μg

Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Background

Human HCC-4, also named NCC-4, liver-expressed chemokine (LEC), and lymphocyte and monocyte chemoattractant (LMC), is a novel CC chemokine identified through bioinformatics. HCC-4 cDNA encodes a 120 amino acid (aa) residue precursor protein with a 23 aa residue predicted signal peptide that is cleaved to generate a 97 aa residue mature protein. HCC-4 is distantly related to other CC chemokines, exhibiting less than 30% aa sequence identity. Among these CC chemokines, HCC-4 has the most similarity to HCC-1. Two potential polyadenylation signals are present on the human HCC-4 gene, and as a result, two transcripts containing approximately 1,500 base pairs and 500 base pairs have been detected. HCC-4 is expressed weakly by some lymphocytes, including NK cells, gamma δ T cells, and some T cell clones. The expression of HCC-4 in monocytes is highly upregulated in the presence of IL-10. The HCC-4 gene has been mapped to chromosome 17q where multiple CC chemokines are clustered. Recombinant HCC-4 has been shown to chemoattract human monocytes and THP-1 cells but not resting lymphocytes or neutrophils. HCC-4 has also been found to suppress proliferation of myeloid progenitor cells. The HCC-4 induced calcium flux in THP-1 cells can be desensitized by prior exposure to RANTES, suggesting that HCC-4 and RANTES share the same receptor in THP-1 cells.

Basic Information

Description


Recombinant Human CCL16/HCC-4 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln24-Gln120) of Human CCL16/HCC-4 (Accession #NP_004581.1) fused with a His tag at the C-terminus.

Bio-Activity

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human CCL16/HCC-4 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 15-20kDa.