

# Recombinant Mouse CCL2/MCP-1 Protein

Catalog No.: RP01626 **Recombinant**

## Sequence Information

Species	Gene ID	Swiss Prot
Mouse	20296	P10148

### Tags

No tag

### Synonyms

C-C motif chemokine ligand 2; CCL2;  
GDCF-2; HC11; HSMCR30; MCAF; Mcp1;  
MCP-1; SCYA2; SMC-CF;CCL2

## Product Information

Source	Purification
<i>Pichia</i>	> 95% by SDS-PAGE.

### Endotoxin

<0.1EU/ $\mu$ g

### Formulation

Lyophilized from a 0.22  $\mu$ m filtered solution of PBS, pH 7.4.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

Monocyte chemoattractant protein 1 (MCP-1), also called CCL2, belongs to a group of CC chemokines located in chromosome 17q11.2. MCP-1 protein interacts with chemokine C-C motif receptor 2 (CCR2) to activate and recruit monocytes, macrophages, CD4+ T cells and immature dendritic cells to the site of infection. The presence of MCP-1 protein in an adequate concentration is important for granuloma formation and M. tuberculosis clearance.

## Basic Information

### Description

Recombinant Mouse CCL2/MCP-1 Protein is produced by *Pichia* expression system. The target protein is expressed with sequence (Gln 24-Asn148) of mouse CCL2/MCP-1 (Accession #NP\_035463.1) fused with no additional amino acid.

### Bio-Activity

1. Measured by the ability to inhibit the proliferation of HUVEC (Human Umbilical Vein Endothelial Cells). The ED<sub>50</sub> for this effect is 0.20-0.80 ng/mL. 2. Measured by its ability to chemoattract THP-1 cells. The ED<sub>50</sub> for this effect is typically 2.87-11.48 ng/mL.


### Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


## Validation Data



Recombinant Mouse CCL2/MCP-1 Protein was determined by SDS-PAGE with Coomassie Blue, showing bands at 14-18 kDa.



Recombinant Mouse CCL2 inhibit the proliferation of HUVEC(Human Umbilical Vein Endothelial Cells).The ED<sub>50</sub> for this effect is 0.20-0.80 ng/mL.



Recombinant Mouse CCL2 chemoattract THP-1 cells. The ED<sub>50</sub> for this effect is typically 2.87-11.48 ng/mL.