Recombinant Human IL-9 Protein



www.abclonal.com

Catalog No.: RP01649 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 3578 P15248

Tags C-hFC

Synonyms

P40; HP40; IL-9;IL-9

Product Information

Source Purification HEK293 cells > 90% by SDS-PAGE.

Endotoxin

<0.1EU/µg

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

a	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Background

IL-9 is a pleiotropic cytokine that influences various distinct functions of different target cells such as T cells, B cells, mast cells and airway epithelial cells by activating STAT1, STAT3 and STAT5. Because of its pleiotropic functions, IL-9 has been demonstrated to be involved in several diseases, such as cancer, autoimmunity and other pathogen-mediated immune-regulated diseases. In this review, we focus on the role of Th9 and IL-9-producing cells in allergic asthma.

Basic Information

Description

Recombinant Human IL-9 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln19-Ile144) of human IL-9 (Accession #) fused with and a hFc tag at the C-terminus.

Bio-Activity

Measured in a cell proliferation assay using MO7e human megakaryocytic leukemic cells. Avanzi, G. et al. (1988) Br. J. Haematol. 69:359. The ED_{50} for this effect is 21.91-87.64 ng/mL.


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human IL-9 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 50-70 kDa.



Recombinant Human IL-9 stimulates cell proliferation of the MO7e human megakaryocytic leukemic cells. The ED $_{50}$ for this effect is 21.91-87.64 ng/mL.