

Recombinant Human IFN-gamma R2 Protein

Catalog No.: RP01650 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 3460 P38484

Tags

C-Rabbit Fc

Synonyms

AF-1; IFGR2; IMD28; IFNGT1; IFNGR2

Product Information

Source Purification HEK293 cells > 97% by SDS-PAGE.

Calculated MW Observed MW

49.71 kDa 65-75 kDa

Endotoxin

<0.1EU/µg

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

a	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Background

Interferon-gamma receptor beta chain (IFNgammaR2), also known as IFNGR2, belongs to the type II cytokine receptor family, whose deficiency is a cause of autosomal recessive mendelian susceptibility to mycobacterial disease (MSMD), also known as familial disseminated atypical mycobacterial infection. This accessory factor is an integral part of the IFN-gamma signal transduction pathway and is likely to interact with GAF, JAK1, and/or JAK2.

Basic Information

Description

Recombinant Human IFN-gamma R2 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ser28-Asp240) of human IFN-gamma R2 (Accession #NP_005525.2) fused with and a RaFc tag at the C-terminus.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20 $^{\circ}$ C for 3 months, at 2-8 $^{\circ}$ C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human IFN-gamma R2 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 50-70 kDa.