Recombinant Human HHG-3/DHH Protein

Catalog No.: RP01758 Recombinant

Sequence Information

| Species | Gene ID | Swiss Prot |
|---------|---------|------------|
| Human | 50846 | 043323 |

Tags

N-6His

Synonyms

GDMN; GDXYM; HHG-3; SRXY7;HHG-3;DHH

Product Information

| Source | Purification | |
|---------|--------------|--|
| E. coli | | |

| Calculated MW | Observed MW |
|---------------|--------------------|
| 20.57 kDa | 24-25 kDa |

Endotoxin

<1 EU/µg

Formulation

Lyophilized from a 0.22 μ m filtered solution of PBS,1mM DTT, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

| 6 | 400-999-6126 |
|-----------|---------------------------|
| \bowtie | cn.market@abclonal.com.cn |
| € | www.abclonal.com.cn |

Background

Desert Hedgehog (Dhh) belongs to the highly conserved Hedgehog family of proteins which are involved in multiple developmental processes. Desert Hedgehog is a secreted, 45 kDa, 373 amino acid (aa) protein that undergoes autocatalytic cleavage between Gly198 and Cys199 catalyzed by the C-terminal domain, which releases the N-terminal domain with a concominant attachment of cholesterol at its new Cterminus. In addition to the C-terminally attached cholesterol, a fatty acid acyl chain is esterified to the N-terminal cysteine (aa 23) via an amide linkage. The 19 kDa Nterminal signaling domain is membrane associated due to its double lipid modifications. Its binding to Patched receptors results in the loss of Patched repression of Smoothened signaling (1 - 4). Dhh binds both Patched and Patched 2 as well as Hedgehog interacting protein (Hip) (5). Within the N-terminal domain, human Dhh shares 97% aa sequence identity with mouse and rat Dhh. It shares approximately 75% aa sequence identity with human Indian (Ihh) and Sonic Hedgehog (Shh) (5). Dhh is produced by Sertoli cells and is required for testis development and spermatogenesis (6, 7). It induces steroidogenic factor 1, which is instrumental in promoting Leydig cell differentiation (8, 9).

Basic Information

Description

Recombinant Human HHG-3/DHH Protein is produced by *E. coli* expression system. The target protein is expressed with sequence (Cys23-Gly198) of human HHG-3/DHH (Accession $\#NP_066382.1$) fused with a 6×His tag at the N-terminus.

Bio-Activity


Storage

Store at -20°C.Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20 $^{\circ}\text{C}$ for 3 months, at 2-8 $^{\circ}\text{C}$ for up to 1 week.

Avoid repeated freeze/thaw cycles.





Recombinant Human HHG-3/DHH Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 20 kDa.