

Recombinant Human BCEI/PS2/TFF1 Protein

Catalog No.: RP01779 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 7031 P04155

Tags

C-His

Synonyms

pS2; BCEI; HPS2; HP1.A; pNR-2; D21S21:TFF1

Product Information

Source HEK293 cells Purification > 97% by SDS-

PAGE.

Endotoxin

 $< 0.001 \, EU/\mu g$

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	Τ	www.abclonal.com.cn

Background

Trefoil Factor 1 (TFF1), also known as pS2, is one of three structurally related secreted proteins that contain trefoil domains. These domains adopt a three-leaved conformation held together by conserved intrachain disulfide bonds. TFF1 is an approximately 7 kDa peptide that plays an important role in epithelial regeneration and wound healing (1). Mature human TFF1 shares 67% amino acid sequence identity with mouse and rat TFF1. It is expressed by goblet cells of the gastric and intestinal mucosa and by conjunctival goblet cells (2-5). TFF1 is a copper-binding protein that can form disulfide-linked homodimers, associate into disulfide-linked complexes with Gastrokine 2, and form non-covalent complexes with the mucin MUC5AC (4, 6-8). Copper enhances TFF1 homodimerization as well as its ability to promote epithelial cell motility, wound healing, and the colonization of H. pylori in stomach and colon epithelia (9, 10). TFF1 is down-regulated during the progression from gastritis to gastric dysplasia to gastric cancer, although it is up-regulated in breast and prostate cancers (11-13). TFF1 inhibits the formation of calcium oxalate crystals, and its excretion in the urine is reduced in patients with kidney stones (14).

Basic Information

Description

Recombinant Human BCEI/PS2/TFF1 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Glu25-Phe84) of human BCEI/PS2/TFF1 (Accession $\#NP_003216.1$) fused with and a $6\times His$ tag at the C-terminus.


Bio-Activity

Storage

Store the lyophilized protein at -20°C to -80°C for 12 months. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human BCEI/PS2/TFF1 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 17 kDa.