

Recombinant Human Neurogranin/NRGN Protein

Catalog No.: RP01814 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 4900 Q92686

Tags

C-hFC

Synonyms

RC3; hng;NRGN;Neurogranin

Product Information

Source Purification HEK293 cells > 97% by SDS-PAGE.

Calculated MW Observed MW

33.45 kDa 45 kDa

Endotoxin

 $< 0.01 EU/\mu g$

Formulation

Lyophilized from a 0.22 µm filtered solution of 20mM Tris 150mM Nacl, pH 7.5.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

Background

Neurogranin is as a small protein encoded by the NRGN gene. it is a calmodulin-binding protein expressed primarily in the brain, particularly in dendritic spines, and participating in the protein kinase C signaling pathway. Neurogranin is the main postsynaptic protein regulating the availability of calmodulin, binding to it in the absence of calcium. Phosphorylation by protein kinase C lowers its binding ability. It is a potential biomarker of neurological and mental diseases.

Basic Information

Description

Recombinant Human Neurogranin/NRGN Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Asp2-Asp78) of human NRGN/Neurogranin (Accession #) fused with hFc tag at the C-terminus.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20 $^{\circ}\text{C}$ for 3 months, at 2-8 $^{\circ}\text{C}$ for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human NRGN/Neurogranin Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 35-45 kDa.