Recombinant Mouse Autotaxin/E-NPP2 Protein

Catalog No.: RP01852 Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
Mouse	18606	Q9R1E6-1

Tags C-His

- -

Synonyms Ectonucleotide

Ectonucleotide pyrophosphatase/phosphodiesterase family member 2; E-NPP 2; 3.1.4.39; Autotaxin; Extracellular lysophospholipase D; LysoPLD[]Enpp2; Npps2; Pdnp2

Product Information

Source Purification HEK293 cells

Calculated MW Observed MW

94.40 kDa 100-130 kDa

Endotoxin

<0.1EU/ μ g of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of 20mM Tris[]150mM NaCl[]pH7.4

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

6	400-999-6126
\times	cn.market@abclonal.com.cn
€	www.abclonal.com.cn

Background

ENPP-2, also known as Autotaxin, belongs to the ectonucleotide pyrophosphatase/phosphodiesterase (NPP) family. Some NPPs hydrolyze phosphates from nucleotides and their derivatives. ENPP-2 shares 40 - 50% identity to ENPP1 & 3, all of which contain a N-terminal intracellular domain, a single transmembrane domain and a large extracellular domain that includes a catalytic domain, two somatomedin-Blike domains, and a C-terminal nuclease-like domain . Unlike ENPP-1 and ENPP-3, ENPP-2 has weak activity against nucleotides, but exhibits a lysophospholipase D activity which allows the formation of lysophosphatidic acid (LPA) and choline from lysophosphatidylcholine . The hydrolysis of nucleotides and lysophospholipids by ENPP-2 is mediated by a single catalytic site . Evidence shows LPA and sphingosine 1phosphate to be specific inhibitors of ENPP-2 . ENPP-2 was originally found to stimulate tumor cell motility and has since been found to enhance tumor invasion and metastasis (and to be up-regulated in several types of carcinomas including breast and lung .

Basic Information

Description

Recombinant Mouse Autotaxin/E-NPP2 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ser49-Ile862) of Mouse Autotaxin/E-NPP2 (Accession #NP_056559.2) fused with a His tag at the C-terminus.

Bio-Activity

Storage

Store at -20°C.Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20 $^{\circ}\text{C}$ for 3 months, at 2-8 $^{\circ}\text{C}$ for up to 1 week.

Avoid repeated freeze/thaw cycles.



180kDa — 140kDa —	=_
100kDa — 75kDa —	==
60kDa —	_
45kDa —	
35kDa —	_
25kDa —	-
15kDa —	-
10kDa —	_

Recombinant Mouse Autotaxin/E-NPP2 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 100-140 kDa.