

# **Recombinant Mouse NCAM-1/CD56 Protein**

Catalog No.: RP01943 Recombinant

# **Sequence Information**

**Species Gene ID Swiss Prot**Mouse 17967 P13595

Tags

C-Avi&His

**Synonyms** 

Ncam1; Ncam; Neural cell adhesion molecule 1; N-CAM-1; NCAM-1; CD56

## **Product Information**

**Source** Purification HEK293 cells ≥ 95 % as

determined by SDS-

PAGE.

Calculated MW Observed MW

79.31 kDa 100-130

**Endotoxin** 

< 0.01 EU/ $\mu$ g of the protein by LAL method

#### **Formulation**

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

#### Reconstitution

Centrifµge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

#### **Contact**

6		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
<u> </u>	П	www.abclonal.com.cn

## **Background**

Neural cell adhesion molecule 1 (NCAM-1: also CD56) is a membrane-bound glycoprotein that plays an important role in nervous system development and function. Mature mouse NCAM-1 consists of a 692 amino acid (aa) extracellular domain (ECD) with five tandem Ig-like domains and two fibronectin type III domains, an 18 aa transmembrane segment, and a 386 aa cytoplasmic domain . Three major splice variants of NCAM-1 are expressed: the 180 kDa full length NCAM-180 isoform, the 140 kDa NCAM-140 isoform which lacks most of the cytoplasmic domain, and the 120 kDa GPI-anchored NCAM-120 isoform that includes the ECD only. Splicing is tissue specific and developmentally regulated . Within the ECD, mouse NCAM-1 shares 94% and 95% aa sequence identity with human and rat NCAM-1, respectively. It is expressed on neurons and glial cells, skeletal muscle, and immune NK cells . NCAM-1 is extensively modified with polysialic acid (PSA) during development, but this addition is decreased in adult tissues . Polysialylation of NCAM-1 is retained in the adult hippocampus where it is important for synaptic plasticity and memory formation . The PSA moiety also participates in the binding of NCAM-1 to heparan sulfate proteoglycans and NCAM-1 mediated migration of olfactory neurons . Proteolytic shedding of NCAM-1 liberates a soluble ECD fragment that can inhibit cortical neurite branching and growth . The NCAM-140 isoform is preferentially expressed on NK cells that robustly secrete cytokines upon activation. Selective up-regulation of the NCAM-140 isoform in a variety of tumors initiates epithelial-mesenchymal transition (EMT) and promotes tumor cell invasion . Finally, NCAM-1 is known to interact with a number of transmembrane and extracellular molecules. NK cell NCAM-1 binds to T cell FGF R1, co-stimulating IL-2 production by T cells. NCAM-1 also forms a noncovalent membrane complex with GFR alpha 1, 2 and 4, generating a receptor for GDNF, NTN and PSP, respectively. And NCAM-1 is reported to form homophilic trans-interactions, and to interact with L1 CAM in cis, and with HSPGs (agrin and collagen XVIII) in trans. In general, these interactions are involved in cell adhesion, migration, and/or process extension.

#### **Basic Information**

#### Description

Recombinant Mouse NCAM-1/CD56 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Leu20-Thr711\_of Mouse NCAM-1/CD56(Accession #P13595) fused with a Avi&His tag at the C-terminus.

#### **Bio-Activity**

#### **Shipping**

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

#### Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80°C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20  $^{\circ}$ C for 3 months, at 2-8  $^{\circ}$ C for up to 1 week.

Avoid repeated freeze/thaw cycles.

# **Operational Notes**

For your safety and health, please wear a lab coat and disposable gloves for handling.

# **Validation Data**



Recombinant Mouse NCAM-1/CD56 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.