# **Recombinant Human CD19 Protein**

ABclonal www.abclonal.com

Catalog No.: RP01972 Recombinant

## **Sequence Information**

Species Gene ID Swiss Prot Human 930 P15391

### **Tags**

C-His

## **Synonyms**

B-lymphocyte antigen CD19; B-lymphocyte surface antigen B4; Differentiation antigen CD19; T-cell surface antigen Leu-12; CD19;

## **Product Information**

Source

**Purification** 

HEK293 cells

Calculated MW Observed MW

29.10 kDa

## **Endotoxin**

< 0.01 EU/µg of the protein by LAL method.

#### **Formulation**

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

## Reconstitution

Centrifµge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

#### **Contact**

<b>a</b>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
$\odot$	www.abclonal.com.cn

## **Background**

B-lymphocyte antigen CD19 is also known as CD19 (Cluster of Differentiation 19), is a single-pass type I membrane protein which contains two Ig-like C2-type (immunoglobulin-like) domains. CD19 is expressed on follicular dendritic cells and B cells. In fact, it is present on B cells from earliest recognizable B-lineage cells during development to B-cell blasts but is lost on maturation to plasma cells. It primarily acts as a B cell co-receptor in conjunction with CD21 and CD81. Upon activation, the cytoplasmic tail of CD19 becomes phosphorylated, which leads to binding by Srcfamily kinases and recruitment of PI-3 kinase. As on T cells, several surface molecules form the antigen receptor and form a complex on B lymphocytes. The (almost) B cell-specific CD19 phosphoglycoprotein is one of these molecules. The others are CD21 and CD81. These surface immunoglobulin (slg)-associated molecules facilitate signal transduction. On living B cells, anti-immunoglobulin antibody mimicking exogenous antigen causes CD19 to b

## **Basic Information**

#### **Description**

Recombinant Human CD19 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Glu21-Arg277) of human CD19 (Accession #NP 001761.3) fused with an 6×His tag at the C-terminus.

#### **Bio-Activity**

#### Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20  $^{\circ}\text{C}$  for 3 months, at 2-8  $^{\circ}\text{C}$  for up to 1 week.

Avoid repeated freeze/thaw cycles.