

Recombinant Human Prostatic Acid Phosphatase /ACPP www.abclonal.com Protein

Catalog No.: RP01976LQ Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 55 P15309

Tags C-6His

Synonyms

ACP3; ACPP; Prostatic acid phosphatase; PAP; EC:3.1.3.2; 5'-nucleotidase; 5'-NT; EC:3.1.3.5; Acid phosphatase 3; Ecto-5'-nucleotidase; Protein tyrosine phosphatase ACP3; EC:3.1.3.48; Thiamine monophosphatase; TMPase; Cleaved into: PAPf39

Product Information

Source

Purification

HEK293 cells ≥ 95 % as determined by SDS-

PAGE.

Calculated MW Observed MW

41.83 KDa 45-50 kDa

Endotoxin

 $< 0.01 \; \text{EU/}\mu\text{g}$ of the protein by LAL method

Formulation

 $0.22~\mu m$ filtered solution of 20mM TrisHCl, 150mM NaCl, pH 7.5.

Reconstitution

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
<u>~</u>	www.abclonal.com.cn

Background

Prostatic acid phosphatase (PAP, or ACPP), also known as Prostatic Acid Phosphatase /ACPP is an enzyme produced by the prostate. As a non-specific phosphomonoesterase, Prostatic acid phosphatase is synthesized and secreted into seminal plasma under androgenic control. The enzyme is a dimer of molecular weight around 100 kDa. Prostatic acid phosphatase is a clinically important protein for its relevance as a biomarker of prostate carcinoma. Furthermore, it has a potential role in fertilization. The major action of PAP is to dephosphorylate macromolecules with the help of catalytic residues (His(12) and Asp(258)) that are located in the cleft between two domains. Cellular prostatic acid phosphatase (cPAcP), an authentic tyrosine phosphatase, is proposed to function as a negative growth regulator of prostate cancer (PCa) cells in part through its dephosphorylation of ErbB-2. cPAcP functions as a neutral protein tyrosine phosphatase (PTP) in prostate cancer cells and dephosphorylates HER-2/ErbB-2/Neu (HER-2: human epidermal growth factor receptor-2) at the phosphotyrosine (p-Tyr) residues. Injection of the secretory isoform of PAP has potent antinociceptive effects in mouse models of chronic pain. This enzyme exhibits ecto-5'-nucleotidase activity, is widely distributed, and implicated in the formation of chronic pain. Additionally, PAP could be a target molecule in specific immunotherapy for patients with nonprostate adenocarcinomas including colon and gastric cancers.

Basic Information

Description

Recombinant Human Prostatic Acid Phosphatase /ACPP Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Lys33-Asp386) of Human Prostatic Acid Phosphatase /ACPP (Accession #NP_001090.2) fused with His at the C-terminus.

Bio-Activity

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.


Storage

Store at -70° C. This product is stable at $\leq -70^{\circ}$ C for up to 1 year from the date of receipt. For optimal storage, aliquot into smaller quantities after centrifugation and store at recommended temperature. Avoid repeated freeze-thaw cycles. Avoid repeated freeze/thaw cycles.

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.

Validation Data



Recombinant Human Prostatic Acid Phosphatase /ACPP Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.