

Biotinylated Recombinant Human TRAIL-R1/CD261 Protein

Catalog No.: RP02083 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	8797	O00220

Tags

C-His&Avi

Synonyms

TNFRSF10A;APO2;CD261;DR4;TRAILR-1;TRAILR1

Product Information

Source	Purification
HEK293 cells	> 95% by Tris-Bis PAGE;> 95% by SEC-HPLC

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Background

Basic Information

Description


Biotinylated Recombinant Human TRAIL R1 / DR4 / TNFRSF10A Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Pro34-Asn239) of Human TRAIL R1 / DR4 / TNFRSF10A fused with His tag and Avi tag at the C-terminal.

Bio-Activity


Storage

Store the lyophilized protein at -20°C to -80°C for long term. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.


Validation Data




Biotinylated Human TRAIL R1 on Tris-Bis PAGE under reduced conditions. The purity is greater than 95%.



The purity of Biotinylated Human TRAIL R1 is greater than 95% as determined by SEC-HPLC.



Immobilized Human TRAIL, No Tag at 5µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Human TRAIL R1, His Tag with the EC₅₀ of 0.31µg/ml determined by ELISA.



Biotinylated Human TRAIL R1, His-Avi Tag immobilized on CM5 Chip can bind Human TRAIL, No Tag with an affinity constant of 0.315 nM as determined in SPR assay (Biacore T200).