

Recombinant Human von Willebrand factor/vWF Protein

Catalog No.: RP02119 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	7450	P04275

Tags

C-His

Synonyms

F8VWF; VWD;VWF;VWD

Product Information

Source	Purification
CHO Cells	> 75 % as determined by SDS-PAGE

Endotoxin

<1EU/μg

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

Von Willebrand Factor (VWF) is a multimeric glycoprotein involved in hemostasis in blood, binds receptors on the surface of platelets and in connective tissue, thereby mediating the adhesion of platelets to sites of vascular injury. From studies it appears that VWF protein uncoils under these circumstances, decelerating passing platelets. VWF protein is deficient or defective in von Willebrand disease (VWD) and is involved in a large number of other diseases, including thrombosis, thrombotic thrombocytopenic purpura, Stroke, Heydes syndrome, possibly hemolytic-uremic syndrome and so on.

Basic Information

Description

Recombinant Human von Willebrand factor/vWF Protein is produced by CHO Cells expression system. The target protein is expressed with sequence (Met1-Lys2813) of human von Willebrand factor/vWF (Accession #AAB59458.1) fused with 6×His tag at the C-terminus.

Bio-Activity


Storage

Store the lyophilized protein at -20°C to -80°C for 12 months.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human von Willebrand factor/vWF Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 260-350 kDa.