

# Recombinant Human TLR4/CD284 Protein

**Catalog No.:** RP02127 **Recombinant**

## Sequence Information

<b>Species</b>	<b>Gene ID</b>	<b>Swiss Prot</b>
Human	7099	O00206-1

### Tags

C-His

### Synonyms

TLR4; ARMD10; CD284; TLR-4; TOLL; toll-like receptor  
4;ARMD10;CD284;TLR-4;TOLL

## Product Information

<b>Source</b>	<b>Purification</b>
Baculovirus-Insect Cells	> 90% by SDS-PAGE.

<b>Calculated MW</b>	<b>Observed MW</b>
70.5 kDa	68 kda

### Endotoxin

&lt; 1.0 EU/μg of the protein by LAL method.

### Formulation

Lyophilized from a 0.22 μm filtered solution of 20mM PB, 500mM NaCl, pH 6.0, 10% glycerol.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

TLR4, also known as TLR-4, is a member of the Toll-like receptor (TLR) family, which plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. They recognize pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents, and mediate the production of cytokines necessary for the development of effective immunity. TLR4 is most abundantly expressed in placenta, and in myelomonocytic subpopulation of the leukocytes. TLR 4 has also been designated as CD284 (cluster of differentiation 284). It has been implicated in signal transduction events induced by lipopolysaccharide (LPS) found in most gram-negative bacteria. TLR4 Cooperates with LY96 and CD14 to mediate the innate immune response to bacterial lipopolysaccharide (LPS). It acts via MYD88, TIRAP and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response. It is also involved in LPS-independent inflammatory responses triggered by Ni(2+).

## Basic Information

### Description

Recombinant Human TLR4/CD284 Protein is produced by Baculovirus-Insect Cells expression system. The target protein is expressed with sequence (Met1-Lys631) of human TLR4/CD284 (Accession #NP\_612564.1) fused with a 6×His tag at the C-terminus.


### Bio-Activity

### Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.

## Validation Data

---



Recombinant Human TLR4/CD284 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 68 kDa.