

Recombinant Human S100-A13 Protein

Catalog No.: RP02188 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	6284	Q99584

Tags

No tag

Synonyms

S100-A13; S100 calcium-binding protein A13; S100A13

Product Information

Source	Purification
<i>E. coli</i>	> 95% by SDS-PAGE.

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

S100A13 is a member of the S100 family of proteins containing 2 EF-hand calcium-binding motifs. It is widely expressed in various types of tissues with a high expression level in thyroid gland. In smooth muscle cells, this protein co-expresses with other family members in the nucleus and in stress fibers, suggesting diverse functions in signal transduction. It plays a role in the export of proteins that lack a signal peptide and are secreted by an alternative pathway. It binds two calcium ions per subunit and one copper ion. Binding of one copper ion does not interfere with calcium binding. It is required for the copper-dependent stress-induced export of IL1A and FGF1. The calcium-free protein binds to lipid vesicles containing phosphatidylserine, but not to vesicles containing phosphatidylcholine.

Basic Information

Description

Recombinant Human S100-A13 Protein is produced by E.coli expression system. The target protein is expressed with sequence (Ala2-Lys98) of human S100A13 (Accession #Q99584).

Bio-Activity


Storage

Store the lyophilized protein at -20°C to -80 °C for long term.

After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human S100-A13 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 5-14 kD.