# **Recombinant Cynomolgus CD3 epsilon/CD3E Protein**

Catalog No.: RP02214 Recombinant

## Sequence Information

Species Gene ID Swiss Prot Cynomolgus 102133065 Q95LI5-1

Tags C-His

#### Synonyms CD3e; CD3E; T3E; FLJ18683; TCRE; CD3-

epsilon; IMD18

# **Product Information**

Source	Purification
HEK293 cells	> 95% as
	determined by Bis-
	Tris PAGE.> 95% as
	determined by HPLC

#### Calculated MW Observed MW 14-16 kDa

11.7 kDa

#### Endotoxin

< 1EU per µg by the LAL method.

#### Formulation

Lyophilized from 0.22um filtered solution in PBS (pH 7.4). Normally 5 % trehalose is added as protectant before lyophilization.

#### Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact

6	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn

### Background

CD3E, is a single-pass type I membrane protein.CD3 (cluster of differentiation 3) T cell co-receptor helps to activate both the cytotoxic T cell (CD8+ naive T cells) and also T helper cells (CD4+ naive T cells). It consists of a protein complex and is composed of four distinct chains. In mammals, the complex contains a CD3y chain, a CD36 chain, and two CD3ɛ chains.

## **Basic Information**

#### Description

Recombinant Cynomolgus CD3 epsilon/CD3E Protein is produced by mammalian expression system. The target protein is expressed with sequence (Gln22-Asp117) of cynomolgus Cynomolgus CD3 Epsilon (Accession #) fused with a His tag at the Cterminus.

#### **Bio-Activity**

Immobilized Cyno CD3 epsilon at 1 µg/mL(100 µL/Well).Dose response curve for Anti-CD3 Ab. with the EC<sub>50</sub> of 0.2  $\mu$ g/mL determined by ELISA.

#### Storage


Store at -20°C.Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.




www.abclonal.com.cn





Cynomolgus CD3E on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

The purity of Cynomolgus CD3E is greater than 95% as determined by SEC-HPLC.



Immobilized Cynomolgus CD3E, His Tag at 0.5 $\mu$ g/ml (100 $\mu$ l/well) on the plate. Dose response curve for Biotinylated Anti-CD3 Antibody, mFc Tag with the EC<sub>50</sub> of 32.7ng/ml determined by ELISA.