

Recombinant Cynomolgus ACE-2 Protein

Catalog No.: RP02267 **Recombinant**

Sequence Information

Species Gene ID Swiss Prot
 Cynomolgus 102130864 A0A2K5X283

Tags

C-His&Avi

Synonyms

ACE-2;ACEH;ACE2; ACE2

Background

Basic Information

Description

Recombinant Cynomolgus ACE-2 Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Gln18-Val739) of Cynomolgus ACE-2 fused with a His tag and Avi at the C-terminal.

Bio-Activity

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.

Product Information

Source

HEK293 cells

Purification

≥ 95 % as determined by Tris-Bis PAGE; ≥ 95 % as determined by HPLC.

Calculated MW

86.5 kDa

Observed MW

95-110 kDa

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.


Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.


Contact

	400-999-6126
	cn.market@abclonal.com.cn
	www.abclonal.com.cn


Validation Data



Recombinant Cynomolgus ACE-2 Protein was determined by Tris-Bis PAGE under reducing conditions.



The purity of Cynomolgus ACE2 is greater than 95% as determined by SEC-HPLC.



Immobilized Cynomolgus ACE2, His Tag at 5 μ g/ml (100 μ l/well) on the plate. Dose response curve for SARS-CoV-2 Spike RBD, hFc Tag with the EC₅₀ of 8.9ng/ml determined by ELISA.